

MATHEMATISCHES FORSCHUNGSINSTITUT OBERWOLFACH

Report No. 31/2016

DOI: 10.4171/OWR/2016/31

Algebraic K-theory and Motivic Cohomology

Organised by

Thomas Geisser, Tokyo

Annette Huber-Klawitter, Freiburg

Uwe Jannsen, Regensburg

Marc Levine, Essen

26 June – 2 July 2016

ABSTRACT. Algebraic K -theory and motivic cohomology have developed together over the last thirty years. Both of these theories rely on a mix of algebraic geometry and homotopy theory for their construction and development, and both have had particularly fruitful applications to problems of algebraic geometry, number theory and quadratic forms. The homotopy-theory aspect has been expanded significantly in recent years with the development of motivic homotopy theory and triangulated categories of motives, and K -theory has provided a guiding light for the development of non-homotopy invariant theories. 19 one-hour talks presented a wide range of latest results on many aspects of the theory and its applications.

Mathematics Subject Classification (2010): 14Cxx, 14Fxx, 19Axx, 19Exx, 19Fxx.

Introduction by the Organisers

Algebraic K -theory and motivic cohomology have developed together over the last thirty years. Both of these theories rely on a mix of algebraic geometry and homotopy theory for their construction and development, and both have had particularly fruitful applications to problems of algebraic geometry, number theory and quadratic forms. The homotopy-theory aspect has been expanded significantly in recent years with the development of motivic homotopy theory and triangulated categories of motives, and K -theory has provided a guiding light for the development of non-homotopy invariant theories.

The workshop program presented a varied series of lectures on the latest developments in the field. The 53 participants came mostly from Europe, but there were large contingents from the USA and Japan, as well as additional participants

from Argentina, Russia and India. The participants ranged from leading experts in the field to younger researchers and also some graduate students. 19 one-hour talks presented a wide range of the latest results on the theory and its applications, reflecting a good mix of nationalities and age groups.

Here is a more detailed description of the talks.

Computations in K -theory and in the homology of linear groups. Hesselholt (with M. Larsen and A. Lindestrauss) constructed a reduced norm map for p -adic K -theory of a division ring over a local field with residue field of characteristic $p > 0$, using the cyclotomic trace map. Morrow extended known results on mod p^n - K -theory to the p -adic case and used this to prove continuity results, infinitesimal versions of weak Lefschetz for the Chow groups and a p -adic version of a conjecture of Kato-Saito. Wendt described his work giving a presentation of the homology of GL_3 of an elliptic curve and its relation to the construction of an elliptic dilogarithm complex. Schlichting described a refinement of Suslin's results on the homology of GL_n to the case of SL_n and E_n .

Categorical constructions. Zakharevich described her approach to the study of $K_0(Var)$ via a categorical scissors congruence construction. Yamazaki presented a framework (developed jointly with Kahn and Saito) for a triangulated category of "motives with modulus", which hopefully will give a good framework for studying non-homotopy invariant phenomena, such as wild ramification. Déglise showed how to construct a category of effective motives over a general base (joint with Bondarko), which admits a reasonable t -structure. Ivorra gave a description of the nearby cycles functor in terms of tubes in non-archimedean geometry. Panin presented aspects of his work with Garkusha on a new description of the motivic stable homotopy category in terms of "framed correspondences" and drew an analogy with Segal's machine for constructing infinite loop spaces.

Theories for topological rings. Tamme presented his work (with Kerz and Saito) describing the "generic fiber" of the comparison map from the K -theory of a smooth proper scheme over a complete DVR its continuous K -theory in terms of a K -theory of the associated rigid analytic space. Using the theory of bornological algebras, Cortiñas showed how to define a theory of rigid cohomology in the non-commutative setting, and gave a description of this in terms of cyclic homology (a joint work with Cuntz, Meyer and Tamme).

Motives and algebraic cycles. Kohrita extended the classical theory of cycles algebraically equivalent to zero and the notion of a universal regular homomorphism, for smooth projective varieties, to the case of smooth varieties and motivic cohomology. He showed the existence of the universal regular homomorphism to a semi-abelian variety for motivic cohomology in a certain range. Tabuada discussed non-commutative versions of Grothendieck's conjecture on the equality of numerical and homological equivalence, and Voevodsky's smash nilpotence conjecture. He showed that the non-commutative and commutative conjectures are equivalent,

gave applications to quadric fibrations and stacks, and showed that under certain conditions, these conjectures are invariant under homological projective duality.

Ancona presented his work proving Grothendieck's conjecture on the positive definiteness of the intersection pairing in a number of new cases. Kahn showed how to interpret the Griffiths group of a smooth projective threefold X (assuming a Chow-Künneth decomposition) as the group of morphisms in the category of motives modulo algebraic equivalence from the Lefschetz motive to the transcendental part of $h_3(X)$. Vial discussed the question: is the field of definition of the intermediate Jacobian (assuming this to be algebraic, or its algebraic part if it is not) of a smooth projective complex variety X the same as the field of definition of X , and showed that this is the case for J_{alg}^3 . In addition, he showed (with Achter and Casalaina-Martin) that the associated Abel-Jacobi map is Galois equivariant.

Arithmetic. Schmidt (with Stix) considered a more general formulation of Grothendieck's section conjecture, involving the entire étale homotopy type rather than just the fundamental group, and obtained a number of "anabelian" statements for varieties that can be embedded as locally closed subschemes of a product of hyperbolic curves. Zhao gave an extension of duality in local class field theory to higher dimensional regular schemes in positive characteristic. Morin described a unified framework and series of conjectures (developed with Flach) for the zeta-values of arithmetic schemes.

Acknowledgement: The MFO and the workshop organizers would like to thank the National Science Foundation for supporting the participation of junior researchers in the workshop by the grant DMS-1049268, "US Junior Oberwolfach Fellows". Moreover, the MFO and the workshop organizers would like to thank the Simons Foundation for supporting Changlong Zhong in the "Simons Visiting Professors" program at the MFO.

Workshop: Algebraic K-theory and Motivic Cohomology**Table of Contents**

Lars Hesselholt (joint with Michael Larsen and Ayelet Lindenstrauss)	
<i>K-theory of division algebras over local fields</i>	1759
Inna Zakharevich	
<i>The annihilator of the Lefschetz motive</i>	1760
Tohru Kohrita	
<i>Algebraic part of motivic cohomology</i>	1762
Matthew Morrow	
<i>Algebraic K-theory and motivic cohomology of formal schemes in characteristic p</i>	1765
Gonçalo Tabuada	
<i>Remarks on Grothendieck's standard conjecture of type D and on Voevodsky's nilpotence conjecture</i>	1768
Giuseppe Ancona	
<i>On the standard conjecture of Hodge type for abelian varieties</i>	1771
Bruno Kahn	
<i>On the Chow motive of a 3-fold</i>	1774
Alexander Schmidt (joint with Jakob Stix)	
<i>Anabelian geometry with étale homotopy types</i>	1775
Takao Yamazaki (joint with Bruno Kahn, Shuji Saito)	
<i>Motives with modulus</i>	1778
Baptiste Morin (joint with Matthias Flach)	
<i>Zeta-values of arithmetic schemes</i>	1780
Guillermo Cortiñas (joint with Joachim Cuntz, Ralf Meyer, Georg Tamme)	
<i>Cyclic homology and rigid cohomology</i>	1783
Charles Vial (joint with Jeffrey Achter and Sebastian Casalaina-Martin)	
<i>On descending cohomology geometrically</i>	1785
Yigeng Zhao	
<i>Étale duality for p-torsion sheaves on semistable schemes over $\mathbb{F}_q[[t]]$</i> ...	1787
Georg Tamme (joint with Moritz Kerz, Shuji Saito)	
<i>K-theory for non-archimedean algebras and spaces</i>	1790
Frédéric Déglise (joint with Mikhail Bondarko)	
<i>Homotopy t-structure and effectivity</i>	1792

Matthias Wendt	
<i>Homology of GL_3 of elliptic curves</i>	1795
Florian Ivorra (joint with Joseph Ayoub and Julien Sebag)	
<i>Motives, nearby cycles and Milnor fibers</i>	1797
Ivan Panin (joint with G.Garkusha)	
<i>A motivic version of Segal's theorem</i>	1800
Marco Schlichting	
<i>Euler class groups and the homology of elementary groups</i>	1801

Abstracts

***K*-theory of division algebras over local fields**

LARS HESSELHOLT

(joint work with Michael Larsen and Ayelet Lindenstrauss)

Suppose that K is a complete discrete valuation field with finite residue field of characteristic p and that D is a central division algebra over K of finite index d . Thirty years ago, Suslin and Yufryakov [3] proved that for all prime numbers $\ell \neq p$ and all positive integers j , there is an isomorphism of ℓ -adic K -groups

$$K_j(D, \mathbb{Z}_\ell) \xrightarrow{\text{Nrd}_{D/K}} K_j(K, \mathbb{Z}_\ell)$$

such that $d\text{Nrd}_{D/K}$ is equal to the norm $N_{D/K}$. We prove the following analogous result for the p -adic K -groups [1].

Theorem. *For every positive integer j , there is an isomorphism*

$$K_j(D, \mathbb{Z}_p) \xrightarrow{\text{Nrd}_{D/K}} K_j(K, \mathbb{Z}_p)$$

with the property that $d\text{Nrd}_{D/K} = N_{D/K}$.

By contrast with the norm $N_{D/K}$ homomorphism, we do not know that the reduced norm isomorphism $\text{Nrd}_{D/K}$ is induced by a map of K -theory spectra. Therefore, we are not able to conclude that a reduced norm isomorphism between the integral K -groups exists; for it is unknown if the p -primary torsion subgroups of the integral K -groups are of bounded exponent.

To prove the ℓ -adic statement, Suslin-Yufryakov use Gabber-Suslin rigidity to reduce the statement to one concerning the ℓ -adic K -groups of the respective residue fields. The additional tool that makes it possible to now prove the p -adic statement is the cyclotomic trace map of Bökstedt-Hsiang-Madsen from K -theory to topological cyclic homology. We let $S \subset K$ be the valuation ring, let $A \subset D$ be the maximal S -order, and consider the cyclotomic trace map

$$K_j(D, \mathbb{Z}_p) \longrightarrow \text{TC}_j(A | D; p, \mathbb{Z}_p)$$

in the case of the exact category of perfect complexes of left A -modules with the chain maps that become quasi-isomorphisms after extension of scalars to D as the weak equivalences; compare [2, Section 1]. It is an isomorphism, for all positive integers j , as is the cyclotomic trace map

$$K_j(K, \mathbb{Z}_p) \longrightarrow \text{TC}_j(S | K; p, \mathbb{Z}_p)$$

The theorem is now proved by showing that, as a graded $\text{TC}_*(S | K; p, \mathbb{Z}_p)$ -module, $\text{TC}_*(A | D; p, \mathbb{Z}_p)$ is free on a single generator in degree 0. We remark that said generator is not in the image of the cyclotomic trace map and that, by comparison, the graded $K_*(K, \mathbb{Z}_p)$ -module $K_*(D, \mathbb{Z}_p)$ is neither free nor finitely generated, except in the trivial case $d = 1$.

REFERENCES

- [1] L. Hesselholt, M. Larsen, and A. Lindenstrauss, *On the K -theory of division algebras over local fields*, in preparation.
- [2] L. Hesselholt and I. Madsen, *On the K -theory of local fields*, Ann. of Math. **158** (2003), 1–113.
- [3] A. A. Suslin and A. V. Yufryakov, *K -theory of local division algebras*, Soviet Math. Dokl. **33** (1986), 794–798.

The annihilator of the Lefschetz motive

INNA ZAKHAREVICH

The Grothendieck ring of varieties is defined to be the free abelian group generated by k -varieties, modulo the relation that for any closed immersion $Y \hookrightarrow X$, we impose the relation that $[X] = [Y] + [X \setminus Y]$; the ring structure is defined by $[X][Y] = [X \times Y]$. In 2014 two longstanding questions about the Grothendieck ring of varieties were answered:

- (1) If two varieties X and Y are piecewise isomorphic then they are equal in the Grothendieck ring; does the converse hold?
- (2) Is the class of the affine line a zero divisor?

Both questions were answered by Borisov, who constructed an element in the kernel of multiplication by the affine line; coincidentally, the proof also constructed two varieties whose classes in the Grothendieck ring are the same but which are not piecewise isomorphic. In this talk we will investigate these questions further by constructing a topological analog of the Grothendieck ring and analyzing its higher homotopy groups. Using this extra structure we will sketch a proof that Borisov's coincidence is not a coincidence at all: that any element in the annihilator of the Lefschetz motive can be represented by a difference of varieties which are equal in the Grothendieck ring but not piecewise isomorphic.

We define an *assembler* in the following manner:

Let \mathcal{C} be a Grothendieck site with an initial object. We denote the full subcategory of noninitial objects by \mathcal{C}° . We say that a family of maps $\{A_i \rightarrow A\}_{i \in I}$ is a *covering family* if it generates a covering sieve in the topology; it is *finite* if I is finite. A covering family is *disjoint* if for any two morphisms $A_i \rightarrow A$ and $A_j \rightarrow A$, the pullback $A_i \times_A A_j$ exists and is equal to the initial object.

An *assembler* is a small Grothendieck site \mathcal{C} satisfying the following extra conditions:

- (I) \mathcal{C} has an initial object \emptyset , and the empty family is a covering family of \emptyset .
- (R) For any A , any two finite disjoint covering families of A have a common refinement which is itself a finite disjoint covering family.
- (M) All morphisms in \mathcal{C} are monomorphisms.

An assembler encodes all of the formal information necessary to keep track of how varieties decompose, while forgetting the particular geometric nature of

varieties. The category of assemblers comes equipped with a functor K to spectra such that for an assembler \mathcal{C} ,

$$\pi_0 K(\mathcal{C}) \cong \text{free abelian group on objects of } \mathcal{C} / \sim,$$

where the equivalence relation says that given a finite disjoint covering family $\{A_i \rightarrow A\}_{i=1}^n$, $[A] = \sum_{i=1}^n [A_i]$ [ZakB]. We write $K_i(\mathcal{C}) = \pi_i K(\mathcal{C})$. Let \mathcal{V}_k be the assembler whose objects are varieties and whose morphisms are locally closed embeddings, $K_0(\mathcal{V}_k)$ is the Grothendieck ring of varieties.

In addition, $K(\mathcal{C})$ preserves further information about the structure of the problem; for example, any element of $K_1(\mathcal{C})$ is given by an object X and two decompositions of X into the same set of pieces X_1, \dots, X_n [ZakC]; in other words, an element of $K_1(\mathcal{C})$ is represented by a piecewise automorphism of X . Thus $K_1(\mathcal{C})$ keeps track of the different ways an object can be decomposed into the same finite set of pieces in two different ways. These higher homotopy groups of $K(\mathcal{C})$ thus preserve a significant amount of data about scissors congruence which can be used to analyze the problems in more detail.

Assemblers also retain sufficient combinatorial data to allow analysis of the homotopy type of $K(\mathcal{C})$; in particular, we have the following localization and dévissage theorems:

Theorem 1 ([ZakB, Theorem C]). *The functor K extends to a functor on simplicial assemblers. For any morphism of simplicial assemblers $g : \mathcal{C} \rightarrow \mathcal{D}$, there exists a simplicial assembler (\mathcal{D}/g) , and a morphism of assemblers $\mathcal{D} \rightarrow (\mathcal{D}/g)$, such that the sequence*

$$K(\mathcal{C}) \rightarrow^g K(\mathcal{D}) \rightarrow K((\mathcal{D}/g))$$

is a cofiber sequence of spectra.

The set of varieties comes with a filtration by the dimension of the variety. This induces a filtration on the Grothendieck ring $K_0(\mathcal{V}_k)$; unfortunately, it is difficult to learn anything directly about $K_0(\mathcal{V}_k)$ from this filtration, since it is prohibitively difficult to compute the associated graded of the filtration. However, this filtration also gives a filtration on the assembler \mathcal{V}_k and therefore a filtration on the K-theory $K(\mathcal{V}_k)$. It is possible to prove the following localization theorem on the assembler of varieties:

Theorem 2 ([ZakA, Theorem A]). *Let $\mathcal{V}_k^{(n)}$ be the assembler of varieties of dimension at most n , and let $\iota_n : \mathcal{V}_k^{(n-1)} \rightarrow \mathcal{V}_k^{(n)}$ be the inclusion of assemblers. Write B_n for the set of birational isomorphism classes of varieties of dimension n . Then the cofiber of the map*

$$K(\iota_n) : K(\mathcal{V}_k^{(n-1)}) \rightarrow K(\mathcal{V}_k^{(n)})$$

is weakly equivalent to $\bigvee_{[X] \in B_n} \Sigma_+^\infty B\text{Aut } k(X)$.

This theorem computes the associated graded of the dimension filtration on $K(\mathcal{V}_k)$. It produces a homologically-graded spectral sequence where the 0-th diagonal converges to the associated graded of the Grothendieck ring of varieties,

starting from groups of the form $\mathbf{Z}[B_n]$. The -1 -st diagonal is 0; the 1-st diagonal contains groups of the form

$$\bigoplus_{[X] \in B_n} (\text{Aut } k(X))^{ab} \oplus \mathbf{Z}/2.$$

This spectral sequence is the connection between the naïve analysis from the beginning of this section and the actual associated graded of the Grothendieck ring of varieties.

The spectral sequence can also be used to analyze the annihilator of the Lefschetz motive $\mathbb{L} = [\mathbb{A}^1]$. This kernel is important because many of the computational techniques that use the Grothendieck ring require it to be localized at \mathbb{L} . Thus any geometric information present in the annihilator of \mathbb{L} is destroyed and can't be detected by these computational techniques. In [Bor], Borisov constructed an element $[X] - [Y]$ in the annihilator of \mathbb{L} , thus showing that this kernel is nontrivial. Surprisingly enough, his proof also showed that $X \times \mathbb{A}^1$ and $Y \times \mathbb{A}^1$ are not birational, thus constructing an element in the kernel of ι_n for some n . Using assemblers we can show that this is not a coincidence:

Theorem 3 ([ZakA, Theorem D]). *Suppose that $\alpha \in K_0(\mathcal{V}_k)$ is in the kernel of multiplication by \mathbb{L} , and write $\alpha = [X] - [Y]$ with X and Y of minimal dimension. Then $X \times \mathbb{A}^1$ and $Y \times \mathbb{A}^1$ are not scissors congruent.*

Thus any minimal representation of an element in the kernel of multiplication by \mathbb{L} gives an element in the kernel of ι_n . The proof of this theorem relies heavily on the higher homotopy groups of $K(\mathcal{V}_k)$, and thus illustrates the principle that the spectrum $K(\mathcal{V}_k)$ contains useful information which is lost in the Grothendieck ring $K_0(\mathcal{V}_k)$.

REFERENCES

- [Bor] Lev Borisov. Class of the affine line is a zero divisor in the Grothendieck ring. <http://arxiv.org/pdf/1412.6194>.
- [ZakA] Inna Zakharevich. The annihilator of the Lefschetz motive. <http://arxiv.org/abs/1506.06200>.
- [ZakB] Inna Zakharevich. The K -theory of assemblers. <http://arxiv.org/abs/1401.3712>.
- [ZakC] Inna Zakharevich. On K_1 of an assembler. <http://arxiv.org/abs/1506.06197>.

Algebraic part of motivic cohomology

TOHRU KOHRITA

For a smooth proper connected scheme X over an algebraically closed field k , the subgroup $A^r(X)$ of the Chow group $CH^r(X)$ consisting of cycles algebraically equivalent to zero is called the *algebraic part* of $CH^r(X)$, and a group homomorphism ϕ from the algebraic part $A^r(X)$ to the group of rational points of an abelian variety A is called a *regular homomorphism* if for any connected smooth proper

scheme T over k pointed at a rational point t_0 , and for any cycle $Y \in CH^r(T \times X)$, the composition

$$T(k) \xrightarrow{w_Y} A^r(X) \xrightarrow{\phi} A(k)$$

is induced by a scheme morphism $T \rightarrow A$ [Sam, Section 2.5]. Here, the map w_Y sends $t \in T(k)$ to $Y(t) - Y(t_0)$, where $Y(t)$ stands for the image of the intersection of cycles $(t \times X) \cdot Y \in CH^{\dim T+r}(T \times X)$ under the proper pushforward along the projection $T \times X \rightarrow X$. A regular homomorphism $\phi : A^r(X) \rightarrow A(k)$ is said *universal* if, given any regular homomorphism $\phi' : A^r(X) \rightarrow A'(k)$, there is a unique homomorphism of abelian varieties $h : A \rightarrow A'$ such that $h \circ \phi = \phi'$ holds. It is classical that the universal regular homomorphism exists for $r = 1$ and $\dim X$. The case $r = 1$ is the theory of Picard varieties and the case $r = \dim X$ coincides with that of Albanese varieties. The existence of universal regular homomorphism for $r = 2$ is also known by Murre [Mur, Theorem A]. It would be worth noting that one of the important ingredients of Murre's proof is the theorem of Merkurjev and Suslin [MS] on norm residue homomorphisms. We also know that universal regular homomorphisms are isomorphisms if $r = 1$, and isomorphisms on torsion if $r = d$ ([Ro, Bl, Mi]) or $r = 2$ ([Mur]).

In this talk, we consider an analogue for arbitrary smooth (not necessarily proper) schemes by replacing Chow groups with motivic cohomology with compact supports. The classical definitions of algebraic part of Chow groups and regular homomorphisms can readily be adapted by using Voevodsky's tensor triangulated category $DM_{Nis}^-(k)$ of motives over k (see [V] and [MVW]). The tensor structure of $DM_{Nis}^-(k)$ is denoted by \otimes , and $M(X)$ (resp., $M^c(X)$) stands for the motive (resp., motive with compact supports) of a scheme X over k .

Given a zero-cycle z on a smooth scheme T over k (or its image in $H_0(T, \mathbb{Z})$) and $Y \in Hom_{DM_{Nis}^-(k)}(M(T) \otimes M^c(X), \mathbb{Z}(n)[m])$, we write the composition

$$M^c(X) \cong \mathbb{Z} \otimes M^c(X) \xrightarrow{z \otimes id_{M^c(X)}} M(T) \otimes M^c(X) \xrightarrow{Y} \mathbb{Z}(n)[m]$$

of morphisms in $DM_{Nis}^-(k)$ as Y_t . Note that Y_t is by definition an element of $H_c^m(X, \mathbb{Z}(n))$.

Definition 1. Let X be a smooth scheme over k . The **algebraic part** of the motivic cohomology group with compact supports $H_c^m(X, \mathbb{Z}(n))$ is defined as

$$H_{c,alg}^m(X, \mathbb{Z}(n)) := \bigcup_{\substack{T, \text{ smooth} \\ \text{connected}}} \text{im}\{H_0(T, \mathbb{Z})^0 \times Hom_{DM_{Nis}^-(k)}(M(T) \otimes M^c(X), \mathbb{Z}(n)[m]) \rightarrow H_c^m(X, \mathbb{Z}(n))\},$$

where the map sends a pair (z, Y) with

$$z \in H_0(T, \mathbb{Z})^0 := \ker\{H_0(T, \mathbb{Z}) \xrightarrow{str_x} H_0(\text{Spec } k, \mathbb{Z})\}$$

and

$$Y \in Hom_{DM_{Nis}^-(k)}(M(T) \otimes M^c(X), \mathbb{Z}(n)[m])$$

to

$$Y_z \in Hom_{DM_{Nis}^-(k)}(M^c(X), \mathbb{Z}(n)[m]) \stackrel{def}{=} H_c^m(X, \mathbb{Z}(n)).$$

Under the assumption of resolution of singularities, it can be shown that for any smooth connected scheme of dimension d over k , the Suslin-Friedlander duality isomorphism ([V, Theorem 4.3.7]) induces an isomorphism between $H_{c,alg}^{2d}(X, \mathbb{Z}(d))$ and $H_0(X, \mathbb{Z})^0$. If X is, in addition, proper and $(m, n) = (2r, r)$ for some integer r , then there is a canonical isomorphism $H_{c,alg}^{2r}(X, \mathbb{Z}(r)) \cong A^r(X)$. The notion of regular homomorphisms also naturally generalizes to this setting.

Definition 2. Let X be a smooth scheme over k and let S be a semi-abelian variety over k . A group homomorphism $\phi : H_{c,alg}^m(X, \mathbb{Z}(n)) \rightarrow S(k)$ is called **regular** if for any smooth connected scheme T over k pointed at $t_0 \in T(k)$ and any $Y \in Hom_{DM_{Nis}^-(k)}(M(T) \otimes M^c(X), \mathbb{Z}(n)[m])$, the composition

$$T(k) \xrightarrow{w_Y} H_{c,alg}^m(X, \mathbb{Z}(n)) \xrightarrow{\phi} S(k)$$

is induced by some scheme morphism $T \rightarrow S$. Here, w_Y sends $t \in T(k)$ to $Y_t - Y_{t_0} \in H_{c,alg}^m(X, \mathbb{Z}(n))$.

A regular homomorphism $\Phi_{c,X}^{m,n} : H_{c,alg}^m(X, \mathbb{Z}(n)) \rightarrow Alg_{c,X}^{m,n}(k)$ is said *universal* if for any regular homomorphism ϕ , there exists a unique homomorphism h of semi-abelian varieties as indicated in the diagram

$$\begin{array}{ccc} H_{c,alg}^m(X, \mathbb{Z}(n)) & \xrightarrow{\Phi_{c,X}^{m,n}} & Alg_{c,X}^{m,n}(k) \\ & \searrow \forall \phi & \downarrow \exists! h \\ & & S(k) \end{array}$$

With this setup, we can prove the following.

Theorem 3 (Existence). Let X be a smooth connected scheme of dimension d over an algebraically closed field k . Then, there exists a universal regular homomorphism

$$\Phi_{c,X}^{m,n} : H_{c,alg}^m(X, \mathbb{Z}(n)) \rightarrow Alg_{c,X}^{m,n}(k)$$

if $m \leq n + 2$ or $(m, n) = (2d, d)$.

Under the assumption of resolution of singularities, one can show that the (Serre’s) Albanese map $alb_X : H_0(X, \mathbb{Z})^0 \rightarrow Alb_X(k)$ constructed in [SS, Ra] agrees with our universal regular homomorphism for $(m, n) = (2d, d)$ via the isomorphism $H_{c,alg}^{2d}(X, \mathbb{Z}(d)) \cong H_0(X, \mathbb{Z})^0$ mentioned right after Definition 1.

For the case $(m, n) = (2, 1)$, we may interpret the universal regular homomorphism for $H_{c,alg}^2(X, \mathbb{Z}(1))$ in terms of the relative Picard group of a smooth compactification \bar{X} of X with a simple normal crossing boundary divisor Z . Below, let us assume that Z is non-empty. Otherwise, it would be the classical proper case, and one would also need to take fppf sheafification in considering a Picard functor below. Let $Pic_{\bar{X}, Z, red}^0$ be the reduction of the identity component of the group scheme representing the functor that sends $T \in Sch/k$ to the relative Picard group $Pic(T \times \bar{X}, T \times Z)$ as in [B-VS]. We can show:

Theorem 4. *Assume resolution of singularities. If X is a smooth connected scheme over k with a smooth compactification \bar{X} with a non-empty simple normal crossing boundary divisor Z , then there is a canonical regular homomorphism*

$$\phi_0 : H_{c,alg}^2(X, \mathbb{Z}(1)) \longrightarrow Pic_{\bar{X},Z,red}^0(k)$$

and it is universal. In particular, $Pic_{\bar{X},Z,red}^0$ depends only on X . Also, ϕ_0 is an isomorphism.

REFERENCES

- [B-VS] Barbieri-Viale, L., Srinivas, V., Albanese and Picard 1-motives., Mémoires de la SMF 87 (2001), vi–104.
- [Bl] Bloch, S., Torsion algebraic cycles and a theorem of Roitman. *Compositio Math.* 39 (1979), no. 1, 107–127.
- [MVW] Mazza, C., Voevodsky, V., Weibel, C., *Lectures on motivic cohomology*, Clay Mathematics Monographs. 2, American Mathematical Society, Providence, RI, 2006.
- [MS] Merkurjev, A., Suslin, A., K -cohomology of Severi-Brauer varieties and norm residue homomorphisms. *Izvestiya Akad. Nauk. SSSR* 46 (1982), English translation: *Math USSR Izvestiya* 21 (1983), 307–340.
- [Mi] Milne, J. S., Zero cycles on algebraic varieties in non-zero characteristic: Roitman’s theorem. *Compositio Math.* 47 (1982), no. 3, 271–287.
- [Mur] Murre, J. P., Application of algebraic K -theory to the theory of algebraic cycles, in: *Proc. Algebraic Geometry Sitges (Barcelona) 1983*, Lecture Notes in Math. 1124, Springer-Verlag, Berlin 1985, 216–261.
- [Ra] Ramachandran, N., Duality of Albanese and Picard 1-motives. Available at: <http://arxiv.org/abs/math/9804042>.
- [Ro] Roitman, A. A., The torsion of the group of 0-cycles modulo rational equivalence. *Ann. of Math.* (2) 111 (1980), no. 3, 553–569.
- [Sam] Samuel, P., Relations d’équivalence en géométrie algébrique. *Proc. Internat. Congress Math.* 1958, 470–487.
- [SS] Spiess, M., Szamuely, T., On the Albanese map for smooth quasi-projective varieties. *Math. Ann.* 325 (2003), no. 1, 1–17.
- [V] Voevodsky, V., Triangulated categories of motives over a field, in: *Cycles, transfers, and motivic homology theories*, *Ann. of Math. Stud.* 143, Princeton Univ. Press, Princeton, NJ, 2000, 188–238.

Algebraic K -theory and motivic cohomology of formal schemes in characteristic p

MATTHEW MORROW

The talk was a survey of the main results of the preprint [6].

1. PRO GEISSER–LEVINE AND BLOCH–KATO–GABBER THEOREM

If A is any \mathbb{F}_p -algebra, then we may consider the natural homomorphisms

$$K_n(A)/p^r \longleftarrow K_n^M(A)/p^r \xrightarrow{\text{dlog}[\cdot]} W_r \Omega_{A,\log}^n,$$

where $W_r \Omega_{A,\log}^n$ (also denoted by $\nu_r^n(A)$ in the literature) is the subgroup of the Hodge–Witt group $W_r \Omega_A^n$ consisting of elements which can be written étale locally as sums of dlog forms, and the map $\text{dlog}[\cdot]$ is given by $\{a_1, \dots, a_n\} \mapsto$

$\mathrm{dlog}[a_1] \cdots \mathrm{dlog}[a_n]$ as usual. If A is regular and local then both of these homomorphisms are known to be isomorphisms: this reduces, via Gersten sequences, to the case that A is a field, in which case the leftwards isomorphism is due to Geisser and Levine [3], who also proved that $K_n(A)$ is p -torsion-free, and the rightwards isomorphism is the Bloch–Kato–Gabber theorem (see [6, Thm. 5.1] for more details and references; also, to avoid issues caused by finite residue fields, we use Kerz–Gabber’s improved Milnor K -theory throughout). Still assuming that A is regular and local, it follows that $K_n(A; \mathbb{Z}/p^r) \cong W_r \Omega_{A, \log}^n$, which satisfactorily calculates the p -adic part of the algebraic K -theory of A .

The focus of the talk was the following analogous calculation for formal schemes:

Theorem 1. *Let A be a regular local \mathbb{F}_p -algebra, and $I \subseteq A$ an ideal; assume that A/I is both F -finite (i.e., finitely generated over its subring of p^{th} -powers) and a “generalised normal crossing divisor” (i.e., whenever Y_1, \dots, Y_N are some of the irreducible components of $\mathrm{Spec} A/I$, then $(\bigcap_i Y_i)_{\mathrm{red}}$ is regular).*

Then the homomorphisms of pro abelian groups

$$\{K_n(A/I^s)/p^r\}_s \longleftarrow \{K_n^M(A/I^s)/p^r\}_s \xrightarrow{\mathrm{dlog}[\cdot]} \{W_r \Omega_{A/I^s, \log}^n\}_s$$

are surjective and have the same kernel, thereby inducing an isomorphism

$$\{K_n(A/I^s)/p^r\}_s \xrightarrow{\cong} \{W_r \Omega_{A/I^s, \log}^n\}_s.$$

Moreover, the pro abelian group $\{K_n(A/I^s)\}_s$ is p -torsion-free.

In other words, although the homomorphisms will not be isomorphisms for A/I^s for any fixed $s > 1$, the obstruction to having an isomorphism $K_n(A/I^s; \mathbb{Z}/p^r) \cong W_r \Omega_{A/I^s, \log}^n$ is Mittag-Leffler zero as $s \rightarrow \infty$.

In the special case that I is principal and A/I is regular, Theorem 1 can be improved: the stated homomorphisms of pro abelian groups are not merely surjective but even isomorphisms. This is likely to be true without these additional assumptions on I and A/I , but certainly necessary results on the p -torsion in the Milnor K -theory of multivariable truncated polynomial algebras remain to be established.

2. APPLICATIONS

Theorem 1 leads to a number of applications concerning K -theory in the presence of infinitesimal thickenings.

2.1. The continuity problem. The continuity problem in algebraic K -theory asks whether the canonical map $K(A) \rightarrow \mathrm{holim}_s K(A/I^s)$ is a weak equivalence after pro-finite completion whenever A is an I -adically complete ring. If A is an \mathbb{F}_p -algebra then it is sufficient to work with the p -adic part of the K -theories, by Gabber rigidity. The first application of Theorem 1 is an affirmative answer to this question in certain cases:

Theorem 2. *With A and I as in Theorem 1, assume moreover that A is I -adically complete. Then the canonical maps*

$$K_n(A; \mathbb{Z}/p^r) \longrightarrow \pi_n \operatorname{holim}_s K_n(A/I^s; \mathbb{Z}/p^r) \longrightarrow \varprojlim_s K_n(A/I^s; \mathbb{Z}/p^r)$$

are isomorphisms for all $n \geq 0$, $r \geq 1$.

In the case in which A/I is regular, this was proved earlier by Geisser and Hesselholt [2].

2.2. The infinitesimal part of the weak Lefschetz conjecture for Chow groups. Let X be a smooth, projective, d -dimensional variety over a perfect field of characteristic p , and $Y \hookrightarrow X$ a smooth ample divisor. Then the Lefschetz conjecture for Chow groups predicts that the restriction map $CH^n(X) \rightarrow CH^n(Y)$ is an isomorphism rationally if $2n < d - 1$. According to the philosophy of Bloch, Esnault, and Kerz proposed in [1], such conjectures can be split into an algebrisation and an infinitesimal part by noting that the map may be rewritten (by the Bloch–Quillen formula) as $H_{\text{Zar}}^i(X, \mathcal{K}_{n,X}) \rightarrow H_{\text{Zar}}^i(Y, \mathcal{K}_{n,Y})$, which factors through $\varprojlim_s H_{\text{Zar}}^i(Y_s, \mathcal{K}_{n,Y_s})$, i.e., the “Chow group of the formal completion of X along Y ”. The second application of Theorem 1 resolves the infinitesimal part of the conjecture:

Theorem 3. *The restriction map $\varprojlim_s H_{\text{Zar}}^i(Y_s, \mathcal{K}_{n,Y_s}) \rightarrow H_{\text{Zar}}^i(Y, \mathcal{K}_{n,Y})$ is an isomorphism rationally if $2n < d - 1$. In fact, whenever $i + n < d - 1$ the restriction map of pro abelian groups*

$$\{H_{\text{Zar}}^i(Y_s, \mathcal{K}_{n,Y_s})\}_s \longrightarrow \{H_{\text{Zar}}^i(Y, \mathcal{K}_{n,Y})\}_s$$

has kernel and cokernel killed by a power of p .

2.3. Kato–Saito’s conjecture. Let X be a smooth, d -dimensional variety over a perfect field of characteristic p . A standard consequence of Gersten’s conjecture (or of the structure of \mathcal{K}_n as a homotopy invariant presheaf with transfer) is that the canonical maps $H_{\text{Zar}}^i(X, \mathcal{K}_n) \rightarrow H_{\text{Nis}}^i(X, \mathcal{K}_{n,\text{Nis}})$ are isomorphisms for all $i, n \geq 0$, and similarly for Milnor K -theory.

Now let $Y \hookrightarrow X$ be a normal crossing divisor. Then it was conjectured by Kato and Saito [5, pg. 256], as part of their higher dimensional class field theory, that the analogous maps

$$\varprojlim_s H_{\text{Zar}}^i(X, \mathcal{K}_{n,(X,Y_s)}^M) \longrightarrow \varprojlim_s H_{\text{Nis}}^i(X, \mathcal{K}_{n,(X,Y_s),\text{Nis}}^M)$$

would also be isomorphisms if the base field were finite and $i = n = d$. The new theory of reciprocity sheaves [4] even predicts that $H_{\text{Zar}}^i(X, \mathcal{K}_{n,(X,Y_s)}^M) \xrightarrow{\cong} H_{\text{Nis}}^i(X, \mathcal{K}_{n,(X,Y_s),\text{Nis}}^M)$ for each fixed $s \geq 1$.

The final application of Theorem 1 which we present here is a p -adic form of Kato–Saito’s conjecture which is valid in all degrees but only established at this time for Quillen K -theory:

Theorem 4. *With X, Y as immediately above, the canonical map of pro abelian groups*

$$\{H_{\text{Zar}}^i(X, \mathcal{K}_{n,(X,Y_s)}/p^r)\}_s \xrightarrow{\cong} \{H_{\text{Nis}}^i(X, \mathcal{K}_{n,(X,Y_s),\text{Nis}}/p^r)\}_s$$

is an isomorphism for all $i, n \geq 0, r \geq 1$.

REFERENCES

- [1] BLOCH, S., ESNAULT, H., AND KERZ, M. p -adic deformation of algebraic cycle classes. *Invent. Math.* 195, 3 (2014), 673–722.
- [2] GEISSER, T., AND HESSELHOLT, L. On the K -theory of complete regular local \mathbb{F}_p -algebras. *Topology* 45, 3 (2006), 475–493.
- [3] GEISSER, T., AND LEVINE, M. The K -theory of fields in characteristic p . *Invent. Math.* 139, 3 (2000), 459–493.
- [4] KAHN, B., SAITO, S., AND YAMAZAKI, T. Reciprocity sheaves, I. *Compositio Math.* (to appear). [arXiv:1402.4201](https://arxiv.org/abs/1402.4201).
- [5] KATO, K., AND SAITO, S. Global class field theory of arithmetic schemes. In *Applications of algebraic K -theory to algebraic geometry and number theory, Part I, II (Boulder, Colo., 1983)*, vol. 55 of *Contemp. Math.* Amer. Math. Soc., Providence, RI, 1986, pp. 255–331.
- [6] MORROW, M. The K -theory and logarithmic Hodge–Witt sheaves of formal schemes in characteristic p . 1512.04703 (2015).

Remarks on Grothendieck’s standard conjecture of type D and on Voevodsky’s nilpotence conjecture

GONÇALO TABUADA

Let k be a base field of characteristic zero. Given a smooth projective k -scheme X , let us denote by $Z^*(X)_{\mathbb{Q}}$ the \mathbb{Q} -vector space of algebraic cycles on X . Following Grothendieck [2] (see also Kleiman [5, 6]), the standard conjecture of type D , denoted by $D(X)$, asserts that $Z^*(X)_{\mathbb{Q}}/\sim_{\text{hom}} = Z^*(X)_{\mathbb{Q}}/\sim_{\text{num}}$, where the homological equivalence relation \sim_{hom} is taken with respect to a classical Weil cohomology theory. Following Voevodsky [17], the nilpotence conjecture, denoted by $V(X)$, asserts that $Z^*(X)_{\mathbb{Q}}/\sim_{\text{nil}} = Z^*(X)_{\mathbb{Q}}/\sim_{\text{num}}$, where \sim_{nil} stands for the smash-nilpotence equivalence relation introduced in *loc. cit.* By construction, we have $V(X) \Rightarrow D(X)$. Thanks to the work of Lieberman [13], the conjecture $D(X)$ holds when $\dim(X) \leq 4$ or when X is an abelian variety. In the same vein, thanks to the work of Kahn-Sebastian [3], Voevodsky [17], and Voisin [18], the conjecture $V(X)$ holds when $\dim(X) \leq 2$ or when X is an abelian 3-fold. Beyond the aforementioned cases, the preceding conjectures remain wide open.

A *differential graded (=dg) category* \mathcal{A} is a category enriched over complexes of k -vector spaces; consult Keller’s ICM survey [4]. Every (dg) k -algebra A gives naturally rise to a dg category with a single object. Another source of examples is provided by schemes since the category of perfect complexes $\text{perf}(X)$ of every quasi-compact quasi-separated k -scheme X admits a canonical dg enhancement¹ $\text{perf}_{\text{dg}}(X)$. Following Kontsevich [7, 8, 9], a dg category \mathcal{A} is called *smooth* if it is compact as a bimodule over itself and proper if $\sum_n \dim H^n \mathcal{A}(x, y) < \infty$ for every

¹When X is quasi-projective this dg enhancement is unique; see Lunts-Orlov [14, Thm. 2.12].

pair of objects (x, y) . Examples include the dg categories $\text{per}_{\text{dg}}(X)$ associated to smooth proper (quasi-compact quasi-separated) k -schemes X .

Given a smooth proper dg category \mathcal{A} , let $K_0(\mathcal{A})_{\mathbb{Q}}$ be the \mathbb{Q} -linearization of the Grothendieck group of \mathcal{A} . Following [1, 16], this \mathbb{Q} -vector space comes equipped with a smash-nilpotence equivalence relation \sim_{nil} , with an homological equivalence relation \sim_{hom} , and also with a numerical equivalence relation \sim_{num} . Motivated by the original conjectures of Grothendieck and Voevodsky, we hence introduce the following (noncommutative) conjectures:

Conjecture $D_{\text{nc}}(\mathcal{A})$: We have $K_0(\mathcal{A})_{\mathbb{Q}}/\sim_{\text{hom}} = K_0(\mathcal{A})_{\mathbb{Q}}/\sim_{\text{num}}$.

Conjecture $V_{\text{nc}}(\mathcal{A})$: We have $K_0(\mathcal{A})_{\mathbb{Q}}/\sim_{\text{nil}} = K_0(\mathcal{A})_{\mathbb{Q}}/\sim_{\text{num}}$.

Similarly to the commutative case, we have $V_{\text{nc}}(\mathcal{A}) \Rightarrow D_{\text{nc}}(\mathcal{A})$. Our first main result, whose proof makes use of the recent theory of noncommutative motives [15], is the following (consult [1, Thm. 1.1] and [16, Thm. 1.1]):

Theorem 1. *Given a smooth projective k -scheme X , we have the equivalences of conjectures $D(X) \Leftrightarrow D_{\text{nc}}(\text{perf}_{\text{dg}}(X))$ and $V(X) \Leftrightarrow V_{\text{nc}}(\text{perf}_{\text{dg}}(X))$.*

Roughly speaking, Theorem 1 extends Grothendieck’s standard conjecture of type D and Voevodsky’s nilpotence conjecture from the realm of schemes to the broad setting of dg categories. This noncommutative viewpoint enables the proof of the original conjectures of Grothendieck and Voevodsky in several (new) cases. Here are some families of examples:

Quadric fibrations. Let S be a smooth projective k -scheme and $q: Q \rightarrow S$ a flat quadric fibration of relative dimension d . Recall from Kuznetsov [11, §3] the construction of the sheaf $\mathcal{Cl}_0(q)$ of even Clifford algebras associated to q . Our second main result is the following (consult [1, Thm. 1.2] and [16, Thm. 1.2]):

Theorem 2. *We have the following equivalences of conjectures:*

$$D(Q) \Leftrightarrow D_{\text{nc}}(\text{perf}_{\text{dg}}(S, \mathcal{Cl}_0(q))) + D(S) \quad V(Q) \Leftrightarrow V_{\text{nc}}(\text{perf}_{\text{dg}}(S, \mathcal{Cl}_0(q))) + V(S).$$

Moreover, when d is even, the discriminant divisor of q is smooth, and $\dim(S) \leq 4$, resp. $\dim(S) \leq 2$, the conjecture $D(Q)$, resp. $V(Q)$, holds.

Homological projective duality. Let V be a finite dimensional k -vector space and X a smooth projective k -scheme equipped with a map $f: X \rightarrow \mathbb{P}(V)$; we write $\mathcal{O}_X(1)$ for the line bundle $f^*\mathcal{O}_{\mathbb{P}(V)}(1)$. Assume that the triangulated category $\text{perf}(X)$ admits a Lefschetz decomposition $\langle \mathbb{A}_0, \mathbb{A}_1(1), \dots, \mathbb{A}_{i-1}(i-1) \rangle$ with respect to $\mathcal{O}_X(1)$ in the sense of Kuznetsov [12, Def. 4.1]. Following [12, Def. 6.1], let $Y \rightarrow \mathbb{P}(V^*)$ be the HPD-dual² of X . Given a generic linear subspace $L \subset V^*$, consider the linear sections $X_L := X \times_{\mathbb{P}(V)} \mathbb{P}(L^\perp)$ and $Y_L := Y \times_{\mathbb{P}(V^*)} \mathbb{P}(L)$. Our third main result is the following (consult [1, Thm. 1.12] and [16, Thm. 1.4]):

Theorem 3. *Let X and Y be as above. Assume that the linear sections X_L and Y_L are smooth, that $\text{codim}(X_L) = \dim(L)$ and $\text{codim}(Y_L) = \dim(L^\perp)$, and that*

²The HPD-dual Y of X is in general a noncommutative variety in the sense of [10, §2.4].

the conjecture $D_{\text{nc}}(\mathbb{A}_{0,\text{dg}})$, resp. $V_{\text{nc}}(\mathbb{A}_{0,\text{dg}})$, holds³. Under these assumptions, we have the equivalence of conjectures $D(X_L) \Leftrightarrow D(Y_L)$, resp. $V(X_L) \Leftrightarrow V(Y_L)$.

Intuitively speaking, Theorem 3 shows that Grothendieck's standard conjecture of type D and Voevodsky's nilpotence conjecture are invariant under homological projective duality. All the assumptions of Theorem 3 are known to hold in the case of linear duality, Veronese-Clifford duality, Grassmannian-Pfaffian duality, spinor duality, Segre-determinantal duality, etc; consult Kuznetsov's ICM survey [10] and the references therein. In the particular case of the Veronese-Clifford duality, we hence conclude, for example, that the intersection of up to five quadric hypersurfaces in an odd-dimensional ambient projective space satisfies Grothendieck's standard conjecture of type D ; consult [1, Thm. 1.4]. In the same vein, the intersection of up to three quadric hypersurfaces in an odd-dimensional ambient projective space satisfies Voevodsky's nilpotence conjecture; consult [16, Thm. 1.7].

Stacks. Theorem 1 allows us to easily extend the conjectures of Grothendieck and Voevodsky from schemes to stacks. Concretely, given a smooth proper Deligne-Mumford stack \mathcal{X} , let $D(\mathcal{X}) := D_{\text{nc}}(\text{perf}_{\text{dg}}(\mathcal{X}))$ and $V(\mathcal{X}) := V_{\text{nc}}(\text{perf}_{\text{dg}}(\mathcal{X}))$. These extended conjectures can be proved, for example, in the case of intersections of bilinear divisors. Let W be a finite dimensional k -vector space, and \mathcal{X} the associated smooth proper Deligne-Mumford stack $(\mathbb{P}(W) \times \mathbb{P}(W))/\mu_2$ equipped with the map $\mathcal{X} \rightarrow \mathbb{P}(S^2W)$, $([w_1], [w_2]) \mapsto [w_1 \otimes w_2 + w_2 \otimes w_1]$. Given a generic linear subspace $L \subset S^2W^*$, the linear section \mathcal{X}_L corresponds to the intersection of the $\dim(L)$ bilinear divisors in \mathcal{X} parametrized by L . Our fourth main result is the following (consult [16, Thm. 1.10]):

Theorem 4. *Assume that $\dim(W)$ is odd and that $\dim(L) \leq 5$, resp. $\dim(L) \leq 3$. Under these assumptions, the conjecture $D(\mathcal{X}_L)$, resp. $V(\mathcal{X}_L)$, holds.*

REFERENCES

- [1] M. Bernardara, M. Marcolli, and G. Tabuada *Some remarks concerning Voevodsky's nilpotence conjecture*. Available at arXiv:1403.0876. To appear in J. Reine Angew. Math.
- [2] A. Grothendieck, *Standard conjectures on algebraic cycles*. 1969 Algebraic Geometry (Internat. Colloq., Tata Inst. Fund. Res., Bombay, 1968) 193–199. Oxford Univ. Press, London.
- [3] B. Kahn and R. Sebastian, *Smash-nilpotent cycles on abelian 3-folds*. Math. Res. Lett. **16** (2009), no. 6, 1007–1010.
- [4] B. Keller, *On differential graded categories*. International Congress of Mathematicians (Madrid), Vol. II, 151–190. Eur. Math. Soc., Zürich (2006).
- [5] S. L. Kleiman, *The standard conjectures*. Motives (Seattle, WA, 1991), 3–20, Proc. Sympos. Pure Math., **55**, Part 1, Amer. Math. Soc., Providence, RI, 1994.
- [6] ———, *Algebraic cycles and the Weil conjectures*. Dix exposés sur la cohomologie des schémas, 359–386, Adv. Stud. Pure Math., **3**, North-Holland, Amsterdam, 1968.
- [7] M. Kontsevich, *Mixed noncommutative motives*. Talk at the Workshop on Homological Mirror Symmetry, Miami, 2010. Available at www-math.mit.edu/auroux/frg/miami10-notes.
- [8] ———, *Notes on motives in finite characteristic*. Algebra, arithmetic, and geometry: in honor of Yu. I. Manin. Vol. II, 213–247, Progr. Math., **270**, Birkhäuser Boston, MA, 2009.

³Here, $\mathbb{A}_{0,\text{dg}}$ stands for the dg enhancement of \mathbb{A}_0 induced from $\text{perf}_{\text{dg}}(X)$.

- [9] ———, *Noncommutative motives*. Talk at the IAS on the occasion of the 61st birthday of Pierre Deligne (2005). Available at <http://video.ias.edu/Geometry-and-Arithmetic>.
- [10] A. Kuznetsov, *Semiorthogonal decompositions in algebraic geometry*. Available at 1404.3143. To appear in Proceedings of the ICM 2014.
- [11] ———, *Derived categories of quadric fibrations and intersections of quadrics*. Adv. Math. **218** (2008), no. 5, 1340–1369.
- [12] ———, *Homological projective duality*. Publ. Math. IHÉS (2007), no. **105**, 157–220.
- [13] D. Lieberman, *Numerical and homological equivalence of algebraic cycles on Hodge manifolds*. Amer. J. Math. **90**, 366–374, 1968.
- [14] V. Lunts and D. Orlov, *Uniqueness of enhancement for triangulated categories*. J. Amer. Math. Soc. **23** (2010), no. 3, 853–908.
- [15] G. Tabuada, *Noncommutative Motives*. With a preface by Yuri I. Manin. University Lecture Series **63**. American Mathematical Society, Providence, RI, 2015.
- [16] ———, *A note on Grothendieck's (noncommutative) standard conjecture of type D*. Available at arXiv:1605.05307.
- [17] V. Voevodsky, *A nilpotence theorem for cycles algebraically equivalent to zero*. Int. Math. Res. Not. IMRN 1995, no. 4, 187–198.
- [18] C. Voisin, *Remarks on zero-cycles of self-products of varieties*. Moduli of vector bundles (Sanda, 1994; Kyoto, 1994), 265–285, LNM **179**, Dekker, New York, 1996.

On the standard conjecture of Hodge type for abelian varieties

GIUSEPPE ANCONA

The aim of this talk is to give some new (and very partial) results on a classical conjecture on algebraic cycles, due to Grothendieck, called standard conjecture of Hodge type. The text is divided in four sections. In the first two we recall the conjecture and give the list of known results. In the last two we state our contribution and give some ideas of its proof.

In what follows k is a base field and (X, L) is a polarized smooth, projective and geometrically connected variety over k of dimension g . By algebraic cycle we will mean an algebraic cycle with rational coefficients.

1. THE CONJECTURE

Consider the algebraic cycles of codimension i whose cohomology class belongs to the primitive cohomology $H_\ell^{2i, \text{prim}}(X)$. Consider on these cycles the numerical equivalence and let $V^i(X)$ be the quotient vector space¹. For $2i \leq g$, we can define a pairing

$$P : V^i(X) \times V^i(X) \rightarrow \mathbb{Q}$$

$$Z_1, Z_2 \mapsto (-1)^i \cdot \deg(Z_1 \cdot Z_2 \cdot L^{g-2i}).$$

The standard conjecture of Hodge type predicts that this pairing is positive definite.

¹A priori this space (as well as the conjecture) depends on ℓ . When $X = A$ is an abelian variety the independency of ℓ is known.

1.1. **Known cases.** When Grothendieck formulated the conjecture he knew the following three facts.

- (1) The conjecture holds in characteristic zero, as a consequence of the Hodge Index Theorem.
- (2) The conjecture holds for surfaces, by Segre, Hodge et al.
- (3) If we know the conjecture for a fiber of a family, then we know it for the generic fiber, in particular the conjecture is reduced to the case when k is finite.

To the author's knowledge the only result in the literature which has been added to this list since then is a theorem of Milne [1].

2. MILNE'S WORK

Definition 1. Define $L^i(X) \subset V^i(X)$ to be the subspace of cycles that can be written as linear combinations of intersections of divisors. These cycles are called Lefschetz classes. Define $E^i(X) \subset V^i(X)$ to be the subspace orthogonal to $L^i(X)$ with respect to the pairing P . These cycles are called exotic classes.

Theorem 1. (Milne [1]) Let $X = A$ be an abelian variety, then the pairing P restricted to $L^i(A)$ is positive definite.

Thanks to Milne's result, in order to fully prove the conjecture for abelian varieties one is reduced to study the pairing P restricted to the space $E^i(A)$.

3. MAIN RESULT

Theorem 2. Suppose that k is finite and $X = A$ is a simple, ordinary, polarized abelian variety. Let K be the biggest totally real field inside $\text{End}(A)_{\mathbb{Q}}$ and \tilde{K} its Galois closure over \mathbb{Q} . If $[\tilde{K} : \mathbb{Q}]$ is odd, then (for all integers i and n) the pairing P restricted to the space $E^i(A^n)$ is not negative definite.

Some comments on this statement.

- Because of 1.1(3), it is very natural to assume k finite.
- Ordinary abelian varieties form an open dense subset of the moduli space of all abelian varieties (of given positive characteristic), so this is also a mild hypothesis.
- Asking that $[\tilde{K} : \mathbb{Q}]$ is odd is a very restrictive hypothesis. It implies in particular that A is of odd dimension.
- It can happen that the space of exotic classes is reduced to zero (in which case the result of Milne suffices for the conjecture). As soon as this space is not reduced to zero our theorem produces a new class on which the pairing P takes positive value.
- There are examples of abelian varieties satisfying our hypothesis and for which there are (or there should be) some exotic classes. In a 1975 letter to Masser, Serre gives an example of a nine dimensional ordinary abelian variety with exotic classes and such that $\text{End}(A)_{\mathbb{Q}} = \mathbb{Q}(\zeta_{19})$, which implies $\tilde{K} = K = \mathbb{Q}(\zeta_{19}) \cap \mathbb{R}$ and $[\tilde{K} : \mathbb{Q}] = 9$.

4. ON THE STRATEGY

4.1. Milne's strategy. Suppose that our abelian variety A lifts to a complex abelian variety \tilde{A} and all divisors also lift. Then Milne's theorem clearly holds because the intersection products one has to compute can be computed on \tilde{A} , where one can use Hodge Index Theorem.

In general one cannot find such a \tilde{A} but (roughly speaking) for each divisor D_i on A one can find, by Honda-Tate, $(\tilde{A}_i, \tilde{D}_i)$ over \mathbb{C} lifting (A, D_i) . In order to put the informations of each \tilde{D}_i together Milne uses a Tannakian argument.

4.2. Some remarks. If one was able to lift also the exotic classes to \mathbb{C} , Milne's argument would go through also in this case. For example, he shows that Hodge conjecture for complex abelian varieties implies the standard conjecture of Hodge type for abelian varieties. Unfortunately lifting exotic classes seems a problem out of reach with the present technology.

Our starting point is a simple remark. Hodge Index Theorem is a cohomological theorem which gives information on all classes of singular cohomology (and not just algebraic ones). Then understanding how far an algebraic class in $H_\ell^{2i}(A)$ is rational (with respect to the rational structure given by the singular cohomology of \tilde{A}) will give informations on the signature of P .

4.3. Our strategy. The idea is to compare two \mathbb{Q} -quadratic forms, q_1 given by P on $V^i(A)$ and the q_2 given by the analogous pairing on the singular cohomology of \tilde{A} . Then one does the following steps.

- (1) Compare $q_1 \otimes \mathbb{Q}_p$ with $q_2 \otimes \mathbb{Q}_p$ (for each p). When the prime is different from the characteristic this will be rather formal. For p equal to the characteristic we will use that A is ordinary.
- (2) Compute $q_2 \otimes \mathbb{R}$. This is done via the Hodge Index Theorem.
- (3) Write the classical product formula on Hilbert symbols for q_1 and q_2 . Thanks to the previous steps all the terms in this product are known except $q_1 \otimes \mathbb{R}$.
- (4) To deduce from the product formula that there is at least a vector on which q_1 is positive one will need the odd hypothesis of the theorem.

REFERENCES

- [1] J. Milne, *Polarizations and Grothendieck's standard conjectures*, Ann. of Math. **155**(2) (2002), 599–610.

On the Chow motive of a 3-fold

BRUNO KAHN

Let X be a smooth projective 3-fold over a field k . Assuming the existence of a Chow-Künneth decomposition for its Chow motive $h(X)$

$$h(X) = \bigoplus_{i=0}^6 h_i(X)$$

we decompose $h_3(X)$ further into two parts

$$h_3(X) = t_3(X) \oplus h_1(J^2)(1)$$

where J^2 is a certain (isogeny class of) abelian variety and $t_3(X)$ is the *transcendental part* of $h_3(X)$. This is a higher-dimensional analogue of the transcendental part of the motive of a surface studied in [2].

The hypothesis on X is satisfied when X is an abelian variety, a complete intersection in projective space or the product of a curve and a surface.

Let Ab^2 be the abelian variety constructed by Murre in [3]: it is the universal abelian variety receiving a regular morphism from algebraically trivial cycles of codimension 2 on X . There is a surjection $\text{Ab}^2 \twoheadrightarrow J^2$, which is an isomorphism under the generalised Hodge or Tate conjecture for $H^3(X)$. (The latter are verified when X is a product of 3 elliptic curves.)

The main result of the talk is an isomorphism

$$(1) \quad \text{Griff}(X) \simeq \mathcal{M}_{\text{alg}}(\mathbb{L}, t_3(X)).$$

Here $\text{Griff}(X)$ is the group of numerically trivial cycles of codimension 2 on X , modulo algebraic equivalence, \mathcal{M}_{alg} is the category of pure motives modulo algebraic equivalence and \mathbb{L} is the Lefschetz motive. (Since we work with \mathbf{Q} coefficients, the above isomorphism is to be taken up to isogeny.)

The Bloch-Beilinson–Murre conjectures on filtrations on Chow groups imply that in fact

$$\mathcal{M}(\mathbb{L}, t_3(X)) \xrightarrow{\sim} \mathcal{M}_{\text{alg}}(\mathbb{L}, t_3(X))$$

where \mathcal{M} is the category of Chow motives. This would imply the following striking fact: up to isogeny, the subgroup of $CH^2(X)$ consisting of cycles algebraically equivalent to 0 is the image of an idempotent algebraic correspondence.

In the special case $X = C \times S$, where C is a curve and S is a surface, (1) yields a surjection (up to isogeny)

$$T(S_{k(C)})/T(S) = \mathcal{M}(h_1(C), t_2(S)) \twoheadrightarrow \mathcal{M}_{\text{alg}}(h_1(C), t_2(S)) = \text{Griff}(C \times S)$$

where $T(S)$ is the Albanese kernel of S . This provides a relationship between the Albanese kernel and the Griffiths group.

Let

$$\begin{array}{ccc} \tilde{S} & \xrightarrow{\pi} & S \\ f \downarrow & & \\ D & & \end{array}$$

be a Lefschetz pencil, with generic fibre Γ . It seems that the methods of [1], combined with the absolute irreducibility of the monodromy on the vanishing cohomology of $H^1(\Gamma)$ imply the following: if k is infinite but finitely generated over its prime field, there is a thin subset $T \subset D(k)$ such that $0 \neq \tilde{\iota}_u \in \text{Griff}(\Gamma_u \times S)$ for $u \notin T$, where $\Gamma_u = f^{-1}(u)$ and $\tilde{\iota}_u$ is the class of the graph of the closed immersion $\iota_u : \Gamma_u \hookrightarrow S$, modified by the Chow-Künneth projectors of Γ_u and S defining $h_1(\Gamma_u)$ and $t_2(S)$.

REFERENCES

- [1] M. Green, P. a. Griffiths, K. Paranjape *Cycles over fields of transcendence degree 1*, Michigan Math. J. **52** (2004), 181–187.
- [2] B. Kahn, J.P. Murre, C. Pedrini *On the transcendental part of the motive of a surface*, in Algebraic cycles and motives (for J.P. Murre’s 75th birthday), LMS Series **344** (2), Cambridge University Press, 2007, 143–202.
- [3] J. P. Murre *Applications of algebraic K-theory to the theory of algebraic cycles*, in Algebraic geometry, Sitges (Barcelona), 1983, 216–261, Lecture Notes in Math. **1124**, Springer 1985.

Anabelian geometry with étale homotopy types

ALEXANDER SCHMIDT

(joint work with Jakob Stix)

Grothendieck’s anabelian philosophy [Gr83] predicts the existence of a class of *anabelian* varieties X that are reconstructible from their étale fundamental group $\pi_1^{\text{ét}}(X, \bar{x})$. All examples of anabelian varieties known so far are of type $K(\pi, 1)$, i.e., their higher étale homotopy groups vanish. For general varieties X , the homotopy theoretic viewpoint suggests to ask the modified question, whether they are reconstructible from their *étale homotopy type* $X_{\text{ét}}$ instead of only $\pi_1^{\text{ét}}(X, \bar{x})$. For varieties X of type $K(\pi, 1)$ this makes no difference since then $X_{\text{ét}}$ is weakly equivalent to the classifying space $B\pi_1^{\text{ét}}(X, \bar{x})$.

Recall that the *étale topological type* $X_{\text{ét}}$ of a scheme X is an object in **pro-ss**, the pro-category of simplicial sets. Any geometric point \bar{x} of X defines a point $\bar{x}_{\text{ét}}$ on $X_{\text{ét}}$. If X is locally noetherian, the fundamental group $\pi_1(X_{\text{ét}}, \bar{x}_{\text{ét}})$ is the usual (in the sense of [SGA3] X §6) étale fundamental group $\pi_1^{\text{ét}}(X, \bar{x})$ and the higher homotopy groups of $X_{\text{ét}}$ are the higher étale homotopy groups of X by definition, cf. [AM69], [Fr82]. Isaksen [Is01] defined a model structure on **pro-ss** and we denote the associated homotopy category by $\text{Ho}(\text{pro-ss})$. When considered as an object of $\text{Ho}(\text{pro-ss})$, we refer to $X_{\text{ét}}$ as the *étale homotopy type* of X . For a pro-simplicial set B , we denote the category of morphisms to B in $\text{Ho}(\text{pro-ss})$ by $\text{Ho}(\text{pro-ss}) \downarrow B$.

For the rest of this talk the letter k always denotes a *finitely generated field extension of \mathbb{Q}* .

In the language of étale homotopy theory, the theorem of Mochizuki and Tamagawa on the anabelian geometry of hyperbolic curves [Mo99, Ta97] can be reformulated as follows.

Theorem 1 ([SS15], Thm. 1.1). *Let X and Y be smooth hyperbolic curves over k . Then the natural map*

$$\mathrm{Isom}_k(X, Y) \longrightarrow \mathrm{Isom}_{\mathrm{Ho}(\mathrm{pro}\text{-}\mathrm{ss})\downarrow k_{\mathrm{et}}}(X_{\mathrm{et}}, Y_{\mathrm{et}})$$

is bijective.

Theorem 2 below constitutes a first step towards a generalisation of Theorem 1 to higher dimensional varieties.

Theorem 2 ([SS15], Thm. 1.2). *Let X and Y be smooth, geometrically connected varieties over k which can be embedded as locally closed subschemes into a product of hyperbolic curves over k . Then the natural map*

$$(*) \quad \mathrm{Isom}_k(X, Y) \longrightarrow \mathrm{Isom}_{\mathrm{Ho}(\mathrm{pro}\text{-}\mathrm{ss})\downarrow k_{\mathrm{et}}}(X_{\mathrm{et}}, Y_{\mathrm{et}})$$

is a split injection with a functorial retraction

$$r : \mathrm{Isom}_{\mathrm{Ho}(\mathrm{pro}\text{-}\mathrm{ss})\downarrow k_{\mathrm{et}}}(X_{\mathrm{et}}, Y_{\mathrm{et}}) \longrightarrow \mathrm{Isom}_k(X, Y).$$

We obtain the following weakly anabelian statement as a trivial corollary.

Corollary 3. *Let X and Y be smooth, geometrically connected varieties over k which can be embedded as locally closed subschemes into a product of hyperbolic curves over k .*

If $X_{\mathrm{et}} \cong Y_{\mathrm{et}}$ in $\mathrm{Ho}(\mathrm{pro}\text{-}\mathrm{ss})\downarrow k_{\mathrm{et}}$, then X and Y are isomorphic as k -varieties.

Remark 4. *By [Is04], the functor $X \mapsto X_{\mathrm{et}}$ from smooth k -schemes to $\mathrm{Ho}(\mathrm{pro}\text{-}\mathrm{ss})\downarrow k_{\mathrm{et}}$ factors through the \mathbb{A}^1 -homotopy category of Morel and Voevodsky [MV99]. In particular, it is not faithful. However, this does not affect Theorem 2 since the schemes occurring there are \mathbb{A}^1 -local.*

In order to investigate whether the map $(*)$ of Theorem 2 is bijective, we may (by Corollary 3) assume that $X = Y$ and consider the kernel of the retraction r . A first piece of information is provided by

Theorem 5 ([SS15], Thm. 1.9). *Let X be a smooth, geometrically connected variety over k which can be embedded as a locally closed subscheme into a product of hyperbolic curves over k . Let γ be in the kernel of the retraction map of Theorem 2:*

$$r : \mathrm{Aut}_{\mathrm{Ho}(\mathrm{pro}\text{-}\mathrm{ss})\downarrow k_{\mathrm{et}}}(X_{\mathrm{et}}) \longrightarrow \mathrm{Aut}_k(X).$$

Then the induced automorphism $\pi_1(\gamma) \in \mathrm{Aut}_{\mathrm{Gal}_k}^{\mathrm{out}}(\pi_1^{\mathrm{et}}(X))$ is class-preserving.

We get more information if X is a *strongly hyperbolic Artin neighbourhood*:

Definition 6. *A strongly hyperbolic Artin neighbourhood is a smooth variety X over k such that there exists a sequence of morphisms*

$$X = X_n \rightarrow X_{n-1} \rightarrow \cdots \rightarrow X_1 \rightarrow X_0 = \mathrm{Spec}(k)$$

such that for all i

- (i) *the morphism $X_i \rightarrow X_{i-1}$ is an elementary fibration into hyperbolic curves, and*
- (ii) *X_i admits an embedding into a product of hyperbolic curves.*

Theorem 7 ([SS15], Thm. 6.2). *Let X be a strongly hyperbolic Artin neighbourhood over k and let $\gamma \in \text{Aut}_{\text{Ho}(\text{pro-ss})\downarrow k_{\text{et}}}(X_{\text{et}})$ be an automorphism with $r(\gamma) = \text{id}_X$. Then $\pi_1(\gamma) \in \text{Aut}_{\text{Gal}_k}^{\text{out}}(\pi_1^{\text{et}}(X))$ is the identity.*

Since strongly hyperbolic Artin-neighbourhoods are of type $K(\pi, 1)$, we can deduce:

Corollary 8. *Let X and Y be strongly hyperbolic Artin neighbourhoods over k . Then the natural map*

$$\text{Isom}_k(X, Y) \longrightarrow \text{Isom}_{\text{Ho}(\text{pro-ss})\downarrow k_{\text{et}}}(X_{\text{et}}, Y_{\text{et}}) = \text{Isom}_{\text{Gal}_k}^{\text{out}}(\pi_1^{\text{et}}(X), \pi_1^{\text{et}}(Y))$$

is bijective.

We remark that by different techniques Hoshi proves in [Ho14] §3 a statement similar to Corollary 8 but restricted to dimension ≤ 4 . Corollary 8 implies the following statement predicted by Grothendieck in his letter to Faltings [Gr83]:

Corollary 9. *Let X be a smooth, geometrically connected variety over k . Then every point of X has a basis of Zariski-neighbourhoods consisting of anabelian varieties, in the sense that k -isomorphisms between any two of these are in bijection with outer G_k -isomorphisms of their respective étale fundamental groups.*

REFERENCES

- [AM69] M. Artin, B. Mazur, *Etale Homotopy*. Lecture Notes in Mathematics **100**, Springer 1969.
- [Fr82] E. M. Friedlander, *Etale homotopy of simplicial schemes*. Annals of Mathematics Studies **104**, Princeton University Press, University of Tokyo Press, 1982, vii+190 pp.
- [Gr83] A. Grothendieck, *Brief an Faltings (27/06/1983)*. in: *Geometric Galois Action 1* (ed. L. Schneps, P. Lochak), LMS Lecture Notes **242**, Cambridge, 1997, 49–58.
- [Ho14] Y. Hoshi, *The Grothendieck conjecture for hyperbolic polycurves of lower dimension*. J. Math. Sci. Univ. Tokyo **21** (2014), no. 2, 153–219.
- [Is01] D. C. Isaksen, *A model structure on the category of pro-simplicial sets*. Trans. Amer. Math. Soc. **353** (2001), no. 7, 2805–2841.
- [Is04] D. C. Isaksen, *Etale realization on the \mathbb{A}^1 -homotopy theory of schemes*. Adv. in Math. **184** (2004), 37–63.
- [Mo99] S. Mochizuki, *The local pro- p anabelian geometry of curves*. Invent. Math. **138** (1999), no. 2, 319–423.
- [MV99] F. Morel, V. Voevodsky, *\mathbb{A}^1 -homotopy theory of schemes*. Publ. Math., Inst. Hautes Étud. Sci. **90** (1999), 45–143.
- [SGA3] M. Demazure, A. Grothendieck, *Schémas en groupes. II: Groupes de type multiplicatif, et structure des schémas en groupes généraux*. Lecture Notes in Mathematics **152**, Springer-Verlag, Berlin-New York, 1970, ix+654 pp. Séminaire de Géométrie Algébrique du Bois Marie 1962–64 (SGA 3), Dirigé par M. Demazure et A. Grothendieck.
- [SS15] A. Schmidt, J. Stix *Anabelian geometry with étale homotopy types*, arXiv:1504.01068 [math.NT].
- [Ta97] A. Tamagawa, *The Grothendieck conjecture for affine curves*. Compos. Math. **109** (1997), no. 2, 135–194.

Motives with modulus

TAKAO YAMAZAKI

(joint work with Bruno Kahn, Shuji Saito)

We report our attempt [2] to generalize Voevodsky’s triangulated category $\mathbf{DM}_{\text{gm}}^{\text{eff}}$ of mixed motives over a field [6] in such a way as to encompass non-homotopy invariant phenomena. Here a traditional notion of “modulus” plays a central role. In the first section, we review how the concept of modulus was used in the theories of generalized Jacobian and Albanese. In the second section, after a brief review of Voevodsky’s $\mathbf{DM}_{\text{gm}}^{\text{eff}}$, we survey our generalization of $\mathbf{DM}_{\text{gm}}^{\text{eff}}$ “with modulus”. Throughout, we work over a perfect field k .

1. MODULUS IN THE THEORY OF GENERALIZED JACOBIAN AND ALBANESE

1.1. Generalized Jacobian. Let C be a smooth proper curve over k , D an effective divisor on it, and x_0 a (fixed) k -rational point on $C \setminus |D|$. The generalized Jacobian $J(C, D)$ of C with modulus D is a commutative algebraic group over k that can be characterized by the universality explained in the next subsection [5]. Its group of k -rational points admits a description

$$J(C, D)(k) \cong \text{Div}^0(C \setminus |D|) / \{\text{div}(f) \mid f \in k(C)_D^\times\},$$

where $k(C)_D^\times := \{f \in k(C)^\times \mid f \equiv 1 \pmod{D}\}$. The following statements hold.

- If $D = 0$, $J(C, D)$ is the classical Jacobian of C (hence an abelian variety).
- If D is reduced, $J(C, D)$ is semi-abelian. The converse holds too.

1.2. Universality. Let G be a commutative algebraic group over k and let $f : C \setminus |D| \rightarrow G$ be a k -morphism. Given $E = \sum_i n_i x_i \in \text{Div}(C \setminus |D|)$, we write $f(E) := \sum_i \text{Tr}_{k(x_i)/k} n_i f(x_i) \in G(k)$. A divisor D' such that $|D'| \subset |D|$ is called a modulus for f if $f(\text{div}(g)) = 0$ holds for any $g \in k(C)_{D'}^\times$. There exists the minimum effective divisor $\text{Mod}(f)$ that is a modulus for f . For example, one has $\text{Mod}(f) = 0$ if f can be extended to a morphism $C \rightarrow G$. Thus, if G is an abelian variety, one always has $\text{Mod}(f) = 0$. Similarly, if G is semi-abelian, $\text{Mod}(f)$ is always reduced. If G is not semi-abelian, $\text{Mod}(f)$ can be non-reduced in general.

Now we can state the universality. The generalized Jacobian $J(C, D)$ comes equipped with a k -morphism $\iota : C \setminus |D| \rightarrow J(C, D)$ such that $\iota(x_0) = 0$ and $\text{Mod}(f) \leq D$. (On rational points, ι simply sends x to the class of $x - x_0$.) Any k -morphism $f : C \setminus |D| \rightarrow G$ with $f(x_0) = 0$ and $\text{Mod}(f) \leq D$ uniquely factors as $f = g \circ \iota$ for some morphism $g : J(C, D) \rightarrow G$ of commutative algebraic groups.

1.3. Generalized Albanese. In a series of papers published in 2008–2013 (see [4] and references therein), Kato and Russell generalized this theory to higher dimensional varieties. Indeed, everything stated in the previous subsection remains true when C is replaced by any smooth projective variety, except that the definition of $\text{Mod}(f)$ should be replaced by a new, much more involved one. Note that for semi-abelian varieties (corresponding to the case of reduced D) this is already established in 1958 by Serre, and of course for abelian varieties (corresponding to

the case $D = 0$) it goes back further to the classical theory of Albanese varieties. Nevertheless a generalization to non-reduced D has emerged only in 21st century.

2. MOTIVES WITH MODULUS

2.1. Voevodsky’s motive and generalized Jacobian. Voevodsky’s definition of $\mathbf{DM}_{\text{gm}}^{\text{eff}}$ is, against expectations, rather simple. We first introduce a category \mathbf{Cor} of finite correspondences. It is an additive category having the same objects as \mathbf{Sm} (the category of smooth varieties) and finite correspondences as morphisms. Then $\mathbf{DM}_{\text{gm}}^{\text{eff}}$ is defined to be the pseudo-abelian envelope of the localization of the homotopy category of bounded complexes $K^b(\mathbf{Cor})$ by two relations arising from Nisnevich Mayer-Vietoris and homotopy invariance. For each $X \in \mathbf{Sm}$, there is an object $M(X) \in \mathbf{DM}_{\text{gm}}^{\text{eff}}$ called the motive of X . Let $X, Y \in \mathbf{Sm}$ and assume that X is proper and equidimensional of dimension d . Then one has

$$\text{Hom}_{\mathbf{DM}_{\text{gm}}^{\text{eff}}}(M(Y), M(X)[-j]) \cong CH^d(X \times Y, j).$$

This is one of the most important properties of Voevodsky’s category $\mathbf{DM}_{\text{gm}}^{\text{eff}}$.

A contravariant functor from \mathbf{Cor} to \mathbf{Ab} that is a sheaf for Nisnevich topology is called a Nisnevich sheaf with transfers. They form an abelian category \mathbf{NST} . The Yoneda functor induces a full faithful functor $i : \mathbf{DM}_{\text{gm}}^{\text{eff}} \rightarrow D(\mathbf{NST})$. A Nisnevich sheaf with transfers F is said to be homotopy invariant if $F(X) \cong F(X \times \mathbb{A}^1)$ for any smooth variety X . For any $K \in \mathbf{DM}_{\text{gm}}^{\text{eff}}$, all homology sheaves of $i(K)$ are homotopy invariant. Now let us consider a smooth proper connected curve C having a k -rational point, and a non-empty effective reduced divisor D on it. Then one has $i(M(C \setminus |D|)) \cong (J(C, D) \oplus \mathbb{Z})[0]$, where $J(C, D)$ is identified with the Nisnevich sheaf with transfers represented by it. Since a commutative algebraic group (regarded as an object of \mathbf{NST}) is homotopy invariant if and only if it is semi-abelian, $J(C, D)$ belongs to $\mathbf{DM}_{\text{gm}}^{\text{eff}}$ if and only if D is reduced.

2.2. Motives with modulus. We construct our “modulus versions” of categories \mathbf{Sm} , \mathbf{Cor} , $\mathbf{DM}_{\text{gm}}^{\text{eff}}$ and \mathbf{NST} , which are denoted by \mathbf{MSm} , \mathbf{MCor} , $\mathbf{MDM}_{\text{gm}}^{\text{eff}}$ and \mathbf{MNST} . An object (X, D) of \mathbf{MSm} is called a modulus pair and consists of a locally integral k -variety X and an effective Cartier divisor D such that $X \setminus |D| \in \mathbf{Sm}$. Starting from this category, we develop construction of categories \mathbf{MCor} , $\mathbf{MDM}_{\text{gm}}^{\text{eff}}$, \mathbf{MNST} and of functors $M : \mathbf{MSm} \rightarrow \mathbf{MDM}_{\text{gm}}^{\text{eff}}$, $i : \mathbf{MDM}_{\text{gm}}^{\text{eff}} \rightarrow D(\mathbf{MNST})$. Roughly speaking it is done by following Voevodsky’s strategy, but largely different in details, for which we refer the readers to [2].

The functor $\omega : \mathbf{MSm} \rightarrow \mathbf{Sm}, \omega(X, D) = X \setminus |D|$ induces $\omega_{\text{gm}} : \mathbf{MDM}_{\text{gm}}^{\text{eff}} \rightarrow \mathbf{DM}_{\text{gm}}^{\text{eff}}$ and $\omega_! : D(\mathbf{MNST}) \rightarrow D(\mathbf{NST})$. They provide us with a very useful relation of our categories with Voevodsky’s. They are also important ingredients in the proof of the following result: Let X be a smooth proper and equidimensional variety of dimension d , and let (Y, E) be any modulus pair. Then one has

$$\text{Hom}_{\mathbf{MDM}_{\text{gm}}^{\text{eff}}}(M(Y, E), M(X, 0)[-j]) \cong CH^d(X \times (Y \setminus |E|), j).$$

It is an important open question to generalize this formula to the case of non-trivial divisor on X . We expect it should be described by Binda-Saito's higher Chow group with modulus [1].

Now let us consider a smooth proper connected curve C having a k -rational point, and a non-reduced effective divisor D on it. We expect that $\omega_i(M(C, D)) \cong (J(C, D) \oplus \mathbb{Z})[0]$. Unfortunately we are unable to prove this statement, but an evidence is given in [3]. A more challenging open problem is to describe $i(M(C, D)) \in D(\mathbf{MNST})$ as a complex of \mathbf{MNST} .

REFERENCES

- [1] Federico Binda, Shuji Saito, *Relative cycles with moduli and regulator maps*. Preprint, arXiv:1412.0385.
- [2] Bruno Kahn, Shuji Saito, Takao Yamazaki, *Motives with modulus*. Preprint, arXiv:1511.07124
- [3] Kay Rülling, Takao Yamazaki, *Suslin homology of relative curves with modulus*. J. Lond. Math. Soc. (2) 93 (2016), no. 3, 567–589.
- [4] Henrik Russell, *Albanese varieties with modulus over a perfect field*. Algebra Number Theory 7 (2013), no. 4, 853–892.
- [5] Jean-Pierre Serre, *Algebraic groups and class fields*. Translated from the French. Graduate Texts in Mathematics, 117. Springer-Verlag, New York, 1988. x+207 pp.
- [6] Vladimir Voevodsky, Andrei Suslin, Eric Friedlander, *Cycles, transfers, and motivic homology theories*. Annals of Mathematics Studies, 143. Princeton University Press, Princeton, NJ, 2000. vi+254 pp.

Zeta-values of arithmetic schemes

BAPTISTE MORIN

(joint work with Matthias Flach)

1. PHILOSOPHY

This talk is based on the preprint [2]. We conjecture that on the category of arithmetic schemes (separated schemes of finite type over the integers), there exist two cohomologies with compact support:

Weil-Arakelov cohomology with compact support: $R\Gamma_{ar,c}(-, A(n))$ with values in the bounded derived category $D^b(\mathbf{LCA})$ of the quasi-abelian category \mathbf{LCA} of locally compact abelian groups [5], for any $A \in \mathbf{LCA}$ of finite ranks and any $n \in \mathbb{Z}$;

Weil-étale cohomology with compact support: $R\Gamma_{W,c}(-, \mathbb{Z}(n))$ with values in the bounded derived category $D^b(\mathbf{Ab})$ of abelian groups, for any $n \in \mathbb{Z}$;

such that the following properties hold.

- (1) $R\Gamma_{W,c}(\mathcal{X}, \mathbb{Z}(n))$ is a perfect complex of abelian groups for any \mathcal{X}, n .
- (2) $R\Gamma_{ar,c}(\mathcal{X}, \mathbb{R}(n))$ is a perfect complex of \mathbb{R} -vector spaces for any \mathcal{X}, n .
- (3) There is a fundamental class $\theta \in H^1(\overline{\text{Spec}(\mathbb{Z})}, \mathbb{R}(0))$ such that

$$\dots \xrightarrow{\cup\theta} H_{ar,c}^i(\mathcal{X}, \mathbb{R}(n)) \xrightarrow{\cup\theta} H_{ar,c}^{i+1}(\mathcal{X}, \mathbb{R}(n)) \xrightarrow{\cup\theta} \dots$$

is an acyclic complex.

- (4) Weil-Arakelov cohomology is exact with respect to coefficients: for example we have an exact triangle

$$R\Gamma_{ar,c}(\mathcal{X}, \mathbb{Z}(n)) \rightarrow R\Gamma_{ar,c}(\mathcal{X}, \mathbb{R}(n)) \rightarrow R\Gamma_{ar,c}(\mathcal{X}, \mathbb{R}/\mathbb{Z}(n)) \rightarrow$$

- (5) There is a natural transformation $R\Gamma_{ar,c}(-, \mathbb{Z}(n)) \rightarrow R\Gamma_{W,c}(-, \mathbb{Z}(n))$ such that, for \mathcal{X}/\mathbb{Z} proper and regular, we have an exact triangle

$$R\Gamma_{dR}(\mathcal{X}_{\mathbb{R}}/\mathbb{R})/F^n[-2] \rightarrow R\Gamma_{ar,c}(\mathcal{X}, \mathbb{Z}(n)) \rightarrow R\Gamma_{W,c}(\mathcal{X}, \mathbb{Z}(n)) \rightarrow$$

- (6) For \mathcal{X}/\mathbb{Z} proper regular of pure dimension d , there is a perfect pairing of locally compact abelian groups

$$H_{ar}^i(\overline{\mathcal{X}}, A(n)) \times H_{ar}^{2d+1-i}(\overline{\mathcal{X}}, A^D(d-n)) \rightarrow H_{ar}^{2d+1}(\overline{\mathcal{X}}, \mathbb{R}/\mathbb{Z}(d)) \rightarrow \mathbb{R}/\mathbb{Z}$$

where $A^D := \underline{\text{Hom}}(A, \mathbb{R}/\mathbb{Z})$ is the Pontryagin dual, and a quasi-isomorphism of perfect complexes of abelian groups

$$R\Gamma_W(\overline{\mathcal{X}}, \mathbb{Z}(n)) \xrightarrow{\sim} R\text{Hom}(R\Gamma_W(\overline{\mathcal{X}}, \mathbb{Z}(d-n)), \mathbb{Z}[-2d-1]).$$

- (7) For \mathcal{X}/\mathbb{Z} proper regular, there is an isomorphism (induced by generalized versions of (5) and (6))

$$R\Gamma_{ar,c}(\mathcal{X}, \mathbb{R}/\mathbb{Z}(n)) \simeq R\Gamma_{W,c}(\mathcal{X}, \mathbb{Z}(n)) \otimes^L \mathbb{R}/\mathbb{Z}.$$

- (8) The vanishing order of the zeta function at $s = n$ is given by

$$\text{ord}_{s=n} \zeta(\mathcal{X}, s) = \sum_{i \in \mathbb{Z}} (-1)^i \cdot i \cdot \dim_{\mathbb{R}} H_{ar,c}^i(\mathcal{X}, \mathbb{R}(n)).$$

- (9) The special value $\zeta^*(\mathcal{X}, n)$ at $s = n$ is given up to sign by

$$\lambda_{\mathcal{X},n}(\zeta^*(\mathcal{X}, n)^{-1} \cdot c(\mathcal{X}, n) \cdot \mathbb{Z}) = \Delta(\mathcal{X}/\mathbb{Z}, n)$$

where $\Delta(\mathcal{X}/\mathbb{Z}, n) := (\det_{\mathbb{Z}} R\Gamma_{W,c}(\mathcal{X}, \mathbb{Z}(n))) \otimes_{\mathbb{Z}} (\det_{\mathbb{Z}} R\Gamma_{dR}(\mathcal{X}/\mathbb{Z})/F^n)$, the isomorphism $\lambda_{\mathcal{X},n} : \mathbb{R} \simeq \Delta(\mathcal{X}/\mathbb{Z}, n) \otimes \mathbb{R}$ is defined below and $c(\mathcal{X}, n) \in \mathbb{Q}^*$ is an explicit non-zero rational number, such that $c(\mathcal{X}, n) = 1$ if $n \leq 0$ and $c(\mathcal{X}, n) = 1$ if \mathcal{X} has characteristic p .

The trivialization $\lambda_{\mathcal{X},n}$ is defined as follows. For $C \in D^b(\text{LCA})$ of finite ranks we consider its tangent complex $T_{\infty}C := R\text{Hom}(R\text{Hom}(C, \mathbb{R}/\mathbb{Z}), \mathbb{R})$. Property (4) above and the fact that T_{∞} is triangulated gives an exact triangle

$$T_{\infty}R\Gamma_{ar,c}(\mathcal{X}, \mathbb{Z}(n)) \rightarrow T_{\infty}R\Gamma_{ar,c}(\mathcal{X}, \mathbb{R}(n)) \rightarrow T_{\infty}R\Gamma_{ar,c}(\mathcal{X}, \mathbb{R}/\mathbb{Z}(n)) \rightarrow$$

By properties (5), (2) and (7), this triangle can be identified with

$$R\Gamma_{dR}(\mathcal{X}_{\mathbb{R}}/\mathbb{R})/F^n[-2] \rightarrow R\Gamma_{ar,c}(\mathcal{X}, \mathbb{R}(n)) \rightarrow R\Gamma_{W,c}(\mathcal{X}, \mathbb{Z}(n))_{\mathbb{R}} \rightarrow$$

The map $\lambda_{\mathcal{X},n}$ is the following composition of isomorphisms:

$$\begin{aligned} \lambda_{\mathcal{X},n} : \mathbb{R} &\xrightarrow{\sim} \det_{\mathbb{R}} R\Gamma_{ar,c}(\mathcal{X}, \mathbb{R}(n)) \\ &\xrightarrow{\sim} (\det_{\mathbb{R}} R\Gamma_{W,c}(\mathcal{X}, \mathbb{Z}(n))_{\mathbb{R}}) \otimes_{\mathbb{R}} (\det_{\mathbb{R}} R\Gamma_{dR}(\mathcal{X}_{\mathbb{R}}/\mathbb{R})/F^n) \xrightarrow{\sim} \Delta(\mathcal{X}/\mathbb{Z}, n)_{\mathbb{R}} \end{aligned}$$

where the first isomorphism is induced by the acyclic complex of Property (3) and the second isomorphism is given by the last exact triangle above. Properties (1)-(9) extend to arbitrary $n \in \mathbb{Z}$ the conjectural picture suggested by Lichtenbaum in [6] for $n = 0$.

We now briefly explain a connexion with Deninger’s program [1]. Recall that C. Deninger conjectured the existence of a cohomology theory $\mathcal{X} \mapsto H_c^i(\mathcal{X}, \mathbb{C})$ on the category of arithmetic schemes, which takes values in ∞ -dimensional \mathbb{C} -vector spaces with an endomorphism Θ , such that for \mathcal{X} of dimension d , we have

$$\zeta(\mathcal{X}, s) = \prod_{i=0}^{i=2d} \det_{\infty} \left(\frac{s - \Theta}{2\pi} \mid H_c^i(\mathcal{X}, \mathbb{C}) \right)^{(-1)^{i+1}}$$

where \det_{∞} is a zeta-regularized determinant. We expect a long exact sequence

$$\dots \rightarrow H_{ar,c}^i(\mathcal{X}, \mathbb{C}(n)) \rightarrow H_c^i(\mathcal{X}, \mathbb{C}) \xrightarrow{\frac{n-\Theta}{2\pi}} H_c^i(\mathcal{X}, \mathbb{C}) \rightarrow H_{ar,c}^{i+1}(\mathcal{X}, \mathbb{C}(n)) \rightarrow \dots$$

such that $\cup\theta : H_{ar,c}^i(\mathcal{X}, \mathbb{C}(n)) \rightarrow H_{ar,c}^{i+1}(\mathcal{X}, \mathbb{C}(n))$ coincides with the composite map

$$H_{ar,c}^i(\mathcal{X}, \mathbb{C}(n)) \rightarrow \text{Ker}(n - \Theta) \hookrightarrow H_c^i(\mathcal{X}, \mathbb{C}) \rightarrow \text{Coker}(n - \Theta) \rightarrow H_{ar,c}^{i+1}(\mathcal{X}, \mathbb{C}(n)).$$

Moreover we expect semi-simplicity in the sense that $\text{Ker}(n - \Theta) \hookrightarrow H_c^i(\mathcal{X}, \mathbb{C}) \rightarrow \text{Coker}(n - \Theta)$ is an isomorphism. This would explain the vanishing order conjecture (8) and would identify $\lambda_{\mathcal{X},n} \otimes \mathbb{C}$ with an isomorphism

$$\mathbb{C} \xrightarrow{\sim} \bigotimes_i \det_{\mathbb{C}}^{(-1)^i} [H_c^i(\mathcal{X}, \mathbb{C}) \xrightarrow{\frac{n-\Theta}{2\pi}} H_c^i(\mathcal{X}, \mathbb{C})] \xrightarrow{\sim} \Delta(\mathcal{X}/\mathbb{Z}, n)_{\mathbb{C}}$$

where the first isomorphism is induced by semi-simplicity.

2. A CONDITIONAL RESULT

Let \mathcal{X}/\mathbb{Z} be a proper regular arithmetic scheme of pure dimension d . For $n \geq 0$, we denote by $\mathbb{Z}(n) := z^n(-, 2n - *)$ Bloch’s cycle complex of sheaves on the étale site \mathcal{X}_{et} . For $n < 0$, we set $\mathbb{Z}(n) := \bigoplus_p j_{p,!} \text{colim } \mu_{p^r}^{\otimes n}[-1]$, where p runs over the set of prime numbers and $j_p : \mathcal{X}[1/p] \rightarrow \mathcal{X}$ is the open immersion. We also denote by $\mathbb{Z}(n)$ the extension of the motivic complex to the Artin-Verdier étale topos $\overline{\mathcal{X}}_{et}$ defined in ([2] App. A). We set $R\Gamma_{dR}(\mathcal{X}/\mathbb{Z})/F^n := R\Gamma(\mathcal{X}_{Zar}, L\Omega_{\mathcal{X}/\mathbb{Z}}^*/F^n)$ where $L\Omega_{\mathcal{X}/\mathbb{Z}}^*$ denotes Illusie’s derived de Rham complex. We need the following

Conjecture AV(\mathcal{X}, n). *There is a perfect pairing of finite groups*

$$H^i(\overline{\mathcal{X}}_{et}, \mathbb{Z}/m(n)) \times H^{2d+1-i}(\overline{\mathcal{X}}_{et}, \mathbb{Z}/m(d-n)) \rightarrow H^{2d+1}(\overline{\mathcal{X}}_{et}, \mathbb{Z}/m(d)) \rightarrow \mathbb{Q}/\mathbb{Z}.$$

Conjecture L(\mathcal{X}, n). *$H^i(\overline{\mathcal{X}}_{et}, \mathbb{Z}(n))$ is finitely generated for any $i \leq 2n + 1$.*

Conjecture B(\mathcal{X}, n). *The pairing*

$$H_c^i(\mathcal{X}, \mathbb{R}(n)) \times H^{2d-i}(\mathcal{X}, \mathbb{R}(d-n)) \rightarrow H_c^{2d}(\mathcal{X}, \mathbb{R}(d)) \rightarrow \mathbb{R}$$

is a perfect pairing of finite dimensional \mathbb{R} -vector spaces. Here $H_c^i(\mathcal{X}, \mathbb{R}(n))$ denotes the cohomology of the mapping fiber of the regulator map.

Theorem. *Let \mathcal{X}/\mathbb{Z} be a proper regular scheme of pure dimension d and let $n \in \mathbb{Z}$. If \mathcal{X} satisfies $\text{AV}(\mathcal{X}, n)$, $\text{L}(\mathcal{X}, n)$, $\text{L}(\mathcal{X}, d - n)$ and $\text{B}(\mathcal{X}, n)$ then there exist Weil-Arakelov complexes $R\Gamma_{ar,c}(\mathcal{X}, A(n))$ and $R\Gamma_{ar}(\overline{\mathcal{X}}, A(n))$ for $A = \mathbb{Z}, \mathbb{R}, \mathbb{R}/\mathbb{Z}$ and Weil-étale complexes $R\Gamma_{W,c}(\mathcal{X}, \mathbb{Z}(n))$ and $R\Gamma_W(\overline{\mathcal{X}}, \mathbb{Z}(n))$, such that:*

- *Properties (1)–(7) hold.*
- *(8) is equivalent to Soulé’s conjecture [7].*
- *If \mathcal{X} has characteristic p , then (9) is equivalent to the Milne-Lichtenbaum-Geisser conjecture [4].*
- *If \mathcal{X} is smooth proper over a number ring \mathcal{O}_F , then (9) is compatible with the Bloch-Kato conjecture [3] for the L -function $L(\oplus_i h^i(\mathcal{X}_F)(n)[-i], s)$.*

In particular, (1)–(8) hold for \mathcal{X}/\mathbb{Z} proper regular of dimension ≤ 1 . If F/\mathbb{Q} is an abelian number field, then (9) holds for $\mathcal{X} = \text{Spec}(\mathcal{O}_F)$ and any $n \in \mathbb{Z}$, and one has $c(\mathcal{X}, n) = (n - 1)!^{-[F:\mathbb{Q}]}$ for $n \geq 1$.

REFERENCES

- [1] C. Deninger, *Some analogies between number theory and dynamical systems on foliated spaces*, Doc. Math. J. DMV. Extra Volume ICM I (1998), 23–46.
- [2] M. Flach, B. Morin, *Weil-étale cohomology and Zeta-values of proper regular arithmetic schemes*, Preprint (2016). arXiv:1605.01277.
- [3] J.M. Fontaine, *Valeurs spéciales des fonctions L des motifs*, Séminaire Bourbaki 751, (1992).
- [4] T. Geisser, *Weil-étale cohomology over finite fields*, Math. Ann. 330 (2004), no. 4, 665–692.
- [5] N. Hoffmann, M. Spitzweck, *Homological algebra with locally compact abelian groups*, Adv. Math. 212 (2007), no. 2, 504–524.
- [6] S. Lichtenbaum, *The Weil-étale Topology for number rings*, Annals of Math. 170 (2009), no. 2, 657–683.
- [7] C. Soulé, *K -théorie et zéros aux points entiers de fonctions zêta*, Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Warsaw, 1983), 437–445.

Cyclic homology and rigid cohomology

GUILLERMO CORTIÑAS

(joint work with Joachim Cuntz, Ralf Meyer, Georg Tamme)

The problem of defining a cohomology theory with good properties for an algebraic variety over a field k of non-zero characteristic has a long history. In the breakthrough paper [8] by Monsky and Washnitzer, such a theory for smooth affine varieties was constructed as follows. Take a complete discrete valuation ring V with uniformizer π and residue field $k = V/\pi V$ (for example, V the Witt ring $W(k)$ if k is perfect). Let K be the fraction field of V . Choose a V -algebra R which is a lift mod π of the coordinate ring of the variety and which is smooth over V (such a lift exists by [4]). Monsky–Washnitzer then introduce the ‘weak’ or dagger-completion R^\dagger of R and define their cohomology as the de Rham cohomology of $R^\dagger \otimes_V K$. The construction of a weak completion has become a basis for the definition of cohomology theories in this context ever since. The

Monsky–Washnitzer theory has been generalized by Berthelot [1] to “rigid cohomology,” which represents a satisfactory cohomology theory for general varieties and schemes over k .

With the advent of cyclic homology around 1980 it was immediately realized that this represents a new approach to de Rham theory. Periodic cyclic homology is defined for arbitrary non-commutative algebras over a field K of characteristic 0, but when specialized to the coordinate ring of a smooth variety (say over \mathbb{C}), it reproduces de Rham theory, see [2],[5], [6]. More generally, it naturally gives the infinitesimal cohomology, in the sense of Grothendieck, for an arbitrary affine variety [5].

Our aim in this project is to develop a version of periodic cyclic theory that reproduces Berthelot’s rigid cohomology for commutative algebras, but which is defined also for non-commutative V -algebras. One starting point is the fact that already in the classical case in characteristic 0 it was found that a natural framework for periodic cyclic homology consists of bornological algebras, see [7]. One of our first results shows that $R^\dagger \otimes_V K$ is a special case of a bornological I -adic completion of $R \otimes_V K$ for an ideal I in R . We use Große-Klönne’s description of Berthelot’s theory to show that in general such bornological I -adic completions can be used very naturally to associate a complete bornological K -algebra B to a given k -algebra A , where K is the quotient field of V , such that the de Rham theory of B gives the rigid cohomology of A . However, to get the analogue of infinitesimal cohomology, we must consider inverse systems representing an ‘infinitesimal neighbourhood’ of an embedding of B into a smooth algebra. This leads to the use of pro-algebras. Accordingly, we have to use the periodic cyclic homology theory developed in [3] for pro-algebras over K , in order to compare periodic cyclic homology for an arbitrary affine variety over k to its rigid cohomology.

Our main result describes rigid cohomology in terms of cyclic homology, as follows. Given a commutative k -algebra A of finite type, write it as a quotient R/J , where R is a smooth (for example, free commutative) V -algebra. Then the rigid cohomology of A is the periodic cyclic homology of the pro-algebra \widetilde{R}_∞ (this notation here is preliminary) defined by the bornological J^m -adic completions of R . We do this by comparing the cyclic homology of \widetilde{R}_∞ to its de Rham cohomology. The de Rham cohomology of \widetilde{R}_∞ recovers the rigid cohomology of A . This description of $H_{\text{rig}}^*(A)$ is in some sense analogous to Grothendieck’s definition of the infinitesimal cohomology of a variety in characteristic 0. However, our definition remains less natural than the description of infinitesimal cohomology via periodic cyclic theory in characteristic 0. This suggests that we may not have the ultimate result.

REFERENCES

- [1] P. Berthelot, *Cohomologie rigide et cohomologie rigide à supports propres. Première partie*, preprint.
- [2] A. Connes, *Noncommutative differential geometry*, Inst. Hautes Études Sci. Publ. Math. **62** (1985), 257–360.

- [3] G. Cortiñas, C. Valqui, *Excision in bivariant periodic cyclic cohomology: a categorical approach*, K-Theory **30** (2003), no. 2, 167–201.
- [4] R. Elkik, *Solutions d'équations à coefficients dans un anneau hensélien*, Ann. Sci. École Norm. Sup. (4) **6** (1973), 553–603.
- [5] B.L. Feigin, B.L. Tsygan, *Additive K-theory and crystalline cohomology*, Funktsional. Anal. i Prilozhen. **19** (1985), no. 2, 52–62.
- [6] J.L. Loday, D. Quillen, *Cyclic homology and the Lie algebra homology of matrices*, Comment. Math. Helv. **59** (1984), no. 4, 569–591.
- [7] R. Meyer, *Local and analytic cyclic homology*, EMS Tracts in Mathematics, vol. 3, European Mathematical Society (EMS), Zürich, 2007.
- [8] P. Monsky, G. Washnitzer, *Formal cohomology. I*, Ann. of Math. (2) **88** (1968), 181–217.

On descending cohomology geometrically

CHARLES VIAL

(joint work with Jeffrey Achter and Sebastian Casalaina-Martin)

Recall that, for a smooth complex projective variety X , the intermediate Jacobian is a complex torus defined by

$$J^{2n+1}(X) := F^{n+1}H^{2n+1}(X, \mathbb{C}) \backslash H^{2n+1}(X, \mathbb{C}) / H^{2n+1}(X, \mathbb{Z}),$$

where F^\bullet denotes the Hodge filtration, and that there is an *Abel–Jacobi map*

$$\text{Ker}(\text{CH}^n(X) \rightarrow H^{2n}(X, \mathbb{Z})) \longrightarrow J^{2n+1}(X).$$

A choice of polarization on X induces a polarization on the Hodge structure $H := H^{2n+1}(X, \mathbb{Z})$. In the case where H has Hodge level 1, *i.e.*, $H \otimes_{\mathbb{Z}} \mathbb{C} = H^{n+1,n} \oplus H^{n,n+1}$, the well-known equivalence of categories between the category of polarized \mathbb{Z} -Hodge structures such that $H \otimes_{\mathbb{Z}} \mathbb{C} = H^{1,0} \oplus H^{0,1}$ and the category of polarized complex abelian varieties shows that $J^{2n+1}(X)$ is in fact a complex abelian variety. At Joe Harris' 60-th birthday conference, Barry Mazur [5] posed the following:

Question 1. Let X be a smooth projective variety defined over a subfield $K \subseteq \mathbb{C}$, and let $X_{\mathbb{C}}$ denote the base-change of X to the field of complex numbers. Assume that $H^{2n+1}(X(\mathbb{C}), \mathbb{Z})$ has Hodge level 1. Does the abelian variety $J^{2n+1}(X_{\mathbb{C}})$ admit a model over K ?

As pointed out by Mazur, this question can be traced back at least to [4], where Deligne gave a positive answer to that question for complete intersections X of odd dimension $2n + 1$ and of Hodge level 1, thereby establishing the Weil conjectures for such X . (Of course, Deligne established the Weil conjectures in full generality not long after.) Deligne proceeded with a proof using the universal family, which consisted in using the irreducibility of the monodromy action on $H^{2n+1}(X(\mathbb{C}), \mathbb{Q})$ and on $H^{2n+1}(X(\mathbb{C}), \mathbb{Z}/\ell)$ for all primes ℓ . It turns out that for such X , one can in fact prove that $H^{2n+1}(X(\mathbb{C}), \mathbb{Q})$ is supported in codimension n , or in other words that $H^{2n+1}(X(\mathbb{C}), \mathbb{Q})$ has geometric coniveau n . The *geometric coniveau filtration*

is defined by :

$$N^\nu H^i(X(\mathbb{C}), \mathbb{Q}) := \sum_{\substack{Z \subseteq X \\ \text{closed, codim} \geq \nu}} \text{Ker} (H^i(X(\mathbb{C}), \mathbb{Q}) \rightarrow H^i((X \setminus Z)(\mathbb{C}), \mathbb{Q})).$$

Resolution of singularities and the formalism of Deligne's mixed Hodge structures show, in particular, that $N^n H^{2n+1}(X(\mathbb{C}), \mathbb{Q})$ is a Hodge structure of Hodge level 1. Note that the generalized Hodge conjecture predicts that $N^n H^{2n+1}(X(\mathbb{C}), \mathbb{Q})$ should be the largest sub-Hodge structure of $H^{2n+1}(X(\mathbb{C}), \mathbb{Q})$ of Hodge level 1. The Hodge structure $N^n H^{2n+1}(X(\mathbb{C}), \mathbb{Q})$ can also be characterized in terms of the Abel–Jacobi map. Define the *algebraic intermediate Jacobian* $J_a^{2n+1}(X_{\mathbb{C}})$ to be the complex torus that is the image of the Abel–Jacobi map restricted to algebraically trivial cycles

$$\text{AJ} : A^{n+1}(X_{\mathbb{C}}) \rightarrow J_a^{2n+1}(X_{\mathbb{C}}).$$

Then $H^1(J_a^{2n+1}(X_{\mathbb{C}}), \mathbb{Q})$ is naturally isomorphic to $N^n H^{2n+1}(X(\mathbb{C}), \mathbb{Q}(n))$ as Hodge structures, and AJ is surjective if and only if $H^{2n+1}(X, \mathbb{Q})$ has geometric coniveau n . One is thus led to ask the following alternative question :

Question 2. Given a smooth projective variety X defined over a subfield $K \subseteq \mathbb{C}$, does the algebraic intermediate Jacobian $J_a^{2n+1}(X_{\mathbb{C}})$ admit a model J over K ? In terms of Galois representations, which was also a motivation for Mazur's question, do there exist an abelian variety J over K and an isomorphism of Galois-representations $N^n H_{\text{ét}}^{2n+1}(X_{\bar{K}}, \mathbb{Q}_{\ell}(n)) \cong H_{\text{ét}}^1(J_{\bar{K}}, \mathbb{Q}_{\ell})$?

In forthcoming work [2], we provide an affirmative answer to Question 2, and as a corollary, give an alternate proof to Deligne's theorem on complete intersections of Hodge level 1.

However, two non-isomorphic abelian varieties over K may become isomorphic after base-change of field. We therefore ask the more precise :

Question 3. Does $J_a^{2n+1}(X_{\mathbb{C}})$ admit a *distinguished* model J over K , in the sense that the Abel–Jacobi map

$$\text{AJ} : A^{n+1}(X_{\mathbb{C}}) \rightarrow J(\mathbb{C})$$

is $\text{Aut}(\mathbb{C}/K)$ -equivariant?

The Picard scheme and the Albanese scheme classically give a positive answer for codimension-1 cycles and for dimension-0 cycles, respectively. In [1], we use Murre's work [6] on algebraic representatives for codimension-2 cycles on smooth projective varieties defined over algebraically closed fields to give a positive answer for codimension-2 cycles :

Theorem. Let X be a smooth projective variety over a field $K \subseteq \mathbb{C}$. The algebraic intermediate Jacobian $J_a^3(X_{\mathbb{C}})$ has a distinguished model J over K . Moreover, there is a correspondence γ on $J \times_K X$ inducing for all primes ℓ a split injective morphism of Galois-representations $H_{\text{ét}}^1(J_{\bar{K}}, \mathbb{Q}_{\ell}) \hookrightarrow H_{\text{ét}}^3(X_{\bar{K}}, \mathbb{Q}_{\ell}(1))$ with image $N^1 H_{\text{ét}}^3(X_{\bar{K}}, \mathbb{Q}_{\ell}(1))$, and with splitting induced by a correspondence γ' on $X \times_K J$.

In categorical terms, we also establish that there exist an abelian variety J over K such that $J_{\mathbb{C}} \cong J_a^{2n+1}(X_{\mathbb{C}})$ and a natural transformation of contravariant functors, extending the Abel–Jacobi map $AJ : A^{n+1}(X_{\mathbb{C}}) \rightarrow J(\mathbb{C})$, from the functor, defined on smooth integral schemes over K , $T \mapsto A^{n+1}(X \times_K T)$ to the functor $T \mapsto J(T)$. An important aspect of our proof, which we will come back to in [3], consists in exhibiting suitable parameter spaces for algebraically trivial cycles. Precisely, we prove and use the following:

Proposition. Let X be a smooth projective variety defined over a perfect field K and consider an algebraically trivial cycle class $\alpha \in A^{n+1}(X_{\bar{K}})$. Then there exist an abelian variety A over K , a cycle class $Z \in \text{CH}^{n+1}(A \times_K X)$, and a pair of \bar{K} -points $t_0, t_1 \in A(\bar{K})$, such that $\alpha = Z_{t_1} - Z_{t_0}$. (Here, Z_{t_i} denotes the Gysin fiber of Z over t_i pushed forward to $X_{\bar{K}}$.)

REFERENCES

- [1] J. Achter, S. Casalaina-Martin, and Ch. Vial, *On descending cohomology geometrically*, preprint.
- [2] J. Achter, S. Casalaina-Martin, and Ch. Vial, *On descending cohomology geometrically, II: descent of the intermediate Jacobian*, forthcoming.
- [3] J. Achter, S. Casalaina-Martin, and Ch. Vial, *Parameter spaces for algebraic equivalence*, forthcoming.
- [4] P. Deligne, *Les intersections complètes de niveau de Hodge un*, Invent. Math. **15** (1972), 237–250.
- [5] B. Mazur, *Open problems: Descending cohomology, geometrically*, Notices of the International Congress of Chinese Mathematicians **2** (2014), no. 1, 37 – 40.
- [6] J. P. Murre, *Applications of algebraic K-theory to the theory of algebraic cycles*, Algebraic geometry, Sitges (Barcelona), 1983, Lecture Notes in Math., vol. 1124, Springer, Berlin, 1985, pp. 216–261.

Étale duality for p -torsion sheaves on semistable schemes over $\mathbb{F}_q[[t]]$

YIGENG ZHAO

1. MOTIVATION

Let K be a local field, i.e., a complete discrete valuation field of characteristic $p > 0$, let \mathcal{O}_K be its ring of integers, let k be its residue field, and let ν_K be its valuation. We fix a uniformizer $\pi \in \mathcal{O}_K$. Recall that in local class field theory, we have the following theorem:

Theorem.(Artin-Schreier-Witt) *There is a perfect pairing of topological groups, that we call the Artin-Schreier-Witt symbol*

$$(1) \quad W_n(K)/(1-F)W_n(K) \times K^\times / (K^\times)^{p^n} \longrightarrow \mathbb{Z}/p^n\mathbb{Z}$$

$$(a, b) \mapsto [a, b] := (b, L/K)(\alpha) - \alpha$$

where $(1-F)(\alpha) = a$, for some $\alpha \in W_n(K^{\text{sep}})$, $L = K(\alpha)$, $(b, L/K)$ is the norm residue of b in L/K , and the topological structure on the first term is discrete, on the second term is induced from K^\times .

Moreover, on both $W_n(K)$ and K^\times , we have filtrations:

$$\begin{aligned} \text{fil}_m W_n(K) &:= \{(a_{n-1}, \dots, a_0) \in W_n(K) \mid p^i \nu_K(a_i) \geq m\}; \\ K^\times \supset \mathcal{O}_K^\times &=: U_K^0 \supset U_K^m := \{x \in \mathcal{O}_K^\times \mid x \cong 1 \pmod{\pi^m}\}. \end{aligned}$$

Theorem. ([1]) *With respect to the Artin-Schreier-Witt symbol (1), the orthogonal complement of $\text{fil}_{m-1} H^1(K, \mathbb{Z}/p^n\mathbb{Z})$ is $U_K^m \cdot (K^\times)^{p^n} / (K^\times)^{p^n}$, for any $m \geq 1$.*

We want to generalize those results to higher dimensional regular schemes.

2. MAIN RESULTS

Let $X \rightarrow \text{Spec}(\mathbb{F}_q[[t]])$ be a projective strictly semistable scheme of relative dimension d , and let $i : X_s \hookrightarrow X$ be its special fiber. Let D be an effective Cartier divisor on X such that $\text{Supp}(D)$ has simple normal crossing, and let $j : U \hookrightarrow X$ be its open complement.

Definition. We define $W_n \Omega_{X|D, \log}^r \subset j_* W_n \Omega_{U, \log}^r$ be the subsheaf étale locally generated by the symbols $\frac{d[x_1]_n}{[x_1]_n} \cdots \frac{d[x_r]_n}{[x_r]_n}$, where $x_1 \in 1 + \mathcal{O}_X(-D)$ and $x_i \in j_* \mathcal{O}_U$ for all i .

If $D_1 \geq D_2$, then $W_n \Omega_{X|D_1, \log}^r \subset W_n \Omega_{X|D_2, \log}^r$. Under this order, we obtain a pro-system “ \varprojlim_D ” $W_m \Omega_{X|D, \log}^r$, where D runs over the subset of effective divisor of X such that $\text{Supp}(D)$ has simple normal crossing and is contained in $X - U$.

Our main theorem is:

Main Theorem. ([4]) *Let $X \rightarrow \text{Spec}(\mathbb{F}_q[[t]])$ be as above. Then there is a perfect pairing of topological $\mathbb{Z}/p^n\mathbb{Z}$ -modules*

$$H^i(U, W_n \Omega_{U, \log}^r) \times \varprojlim_D H_{X_s}^{d+2-i}(X, W_n \Omega_{X|D, \log}^{d+1-r}) \rightarrow H_{X_s}^{d+2}(X, W_n \Omega_{X, \log}^{d+1}) \xrightarrow{\text{Tr}} \mathbb{Z}/p^n\mathbb{Z},$$

where the first term is endowed with the discrete topology, and the second term is endowed with the profinite topology.

The trace morphism in the above theorem is given by the following purity result and Moser-Sato trace map [2, 3] for $\nu_{n, X_s}^d := \text{Ker}(\oplus_{x \in X_s^0} i_x^* W_n \Omega_{x, \log}^d \xrightarrow{\partial} \oplus_{x \in X_s^1} i_x^* W_n \Omega_{x, \log}^{d-1})$, where ∂ is the Kato residue map on logarithmic de Rham-Witt sheaves.

Theorem(Purity). *There is a canonical isomorphism*

$$\text{Gys}_{i, n}^{\log} : \nu_{n, X_s}^d[-1] \xrightarrow{\cong} \text{Ri}^1 W_n \Omega_{X, \log}^{d+1}$$

in $D^+(X_s, \mathbb{Z}/p^n\mathbb{Z})$.

We denote this pairing in our Main Theorem as $\langle \cdot, \cdot \rangle$.

For any $\chi \in H^i(U, W_n \Omega_{U, \log}^r)$, we define the higher Artin conductor

$$\text{ar}(\chi) := \min\{D \mid \langle \chi, \cdot \rangle \text{ factors through } H_{X_s}^{d+2-i}(X, W_n \Omega_{X|D, \log}^{d+1-r})\},$$

Then we define

$$\mathrm{Fil}_D H^i(U, W_n \Omega_{U, \log}^r) := \{\chi \in H^i(U, W_n \Omega_{U, \log}^r) \mid \mathrm{ar}(\chi) \leq D\},$$

and

$$\pi_1^{\mathrm{ab}}(X, D)/p^n := \mathrm{Hom}(\mathrm{Fil}_D H^1(U, \mathbb{Z}/p^n \mathbb{Z}), \mathbb{Q}/\mathbb{Z})$$

endowed with the usual profinite topology of the dual.

The quotient $\pi_1^{\mathrm{ab}}(X, D)/p^n$ classifies abelian étale coverings of U of degree p^n with ramification bounded by the divisor D .

3. APPLICATION

Let $X = B = \mathrm{Spec} \mathbb{F}_q[[t]]$, let $D = X_s = (t)$ be the unique closed point. Then $U = \mathrm{Spec}(\mathbb{F}_q((t)))$. We denote $mD := m(t)$ for any $m \in \mathbb{N}$. Our Main Theorem in this setting is:

$$H^i(K, W_n \Omega_{K, \log}^r) \times \varprojlim_m H_{X_s}^{2-i}(B, W_n \Omega_{B|mD, \log}^{1-j}) \rightarrow \mathbb{Z}/p^n \mathbb{Z}$$

is a perfect pairing of topological abelian groups.

In particular, if $i = 1, j = 0$. We are back to the Artin-Schreier-Witt symbol.

Proposition. *For $m \geq 1$, we have*

$$\mathrm{Fil}_{mD} H^1(K, \mathbb{Z}/p^n \mathbb{Z}) = \mathrm{fil}_{m-1} H^1(K, \mathbb{Z}/p^n \mathbb{Z}),$$

i.e., Our new filtration is same as the Brylinski-Kato filtration.

REFERENCES

- [1] J.-L. Brylinski, *Théorie du corps de classes de Kato et revêtements abéliens de surfaces*, Annales de l'institut Fourier **33** (1983), no. 3, 23–38.
- [2] T. Moser, *A duality theorem for étale p -torsion sheaves on complete varieties over a finite field*, Compositio Mathematica **117** (1999), 123–152.
- [3] K. Sato, *Logarithmic Hodge-Witt sheaves on normal crossing varieties*, Math. Z. **257** (2007), 707–743.
- [4] Y. Zhao, *Étale duality of semistable schemes over local rings of positive characteristic*, Dissertation(2016), Universität Regensburg.

K-theory for non-archimedean algebras and spaces

GEORG TAMME

(joint work with Moritz Kerz, Shuji Saito)

Let K be a complete discretely valued field with ring of integers \mathcal{O}_K , uniformizer π , and residue field k .

1. MOTIVATION

Let $\mathcal{X}/\mathcal{O}_K$ be a smooth proper scheme. Denote by $\mathcal{X}_n := \mathcal{X} \otimes_{\mathcal{O}_K} \mathcal{O}_K/(\pi^n)$ the infinitesimal thickenings of the special fibre \mathcal{X}_k , by $X := \mathcal{X} \otimes_{\mathcal{O}_K} K$ the generic fibre of \mathcal{X} , and by X^{rig} its associated rigid analytic space. By definition, the continuous K -theory of \mathcal{X} is the pro-spectrum $K^{\text{cont}}(\mathcal{X}) := \{K(\mathcal{X}_n)\}_{n \in \mathbb{N}} \in \text{Pro}(\text{Sp})$. The study of the comparison map $K(\mathcal{X}) \rightarrow K^{\text{cont}}(\mathcal{X})$ plays an important role in the strategy to attack Grothendieck's variational Hodge conjecture and its variants proposed by Bloch, Esnault, and Kerz [1, 2]. Our goal here is to understand 'the generic fibre' of this comparison map. More precisely, we want to construct a version of K -theory for the rigid space X^{rig} which fits in a homotopy cartesian square

$$\begin{array}{ccc} K(\mathcal{X}) & \longrightarrow & K^{\text{cont}}(\mathcal{X}) \\ \downarrow & & \downarrow \\ K(X) & \longrightarrow & K\text{-theory of } X^{\text{rig}}. \end{array}$$

2. CONSTRUCTIONS

We first define a K -theory for affinoid algebras, and then globalize it to all rigid spaces. Let A be an affinoid K -algebra. For an integer j define

$$A\langle x \rangle_{\pi^j} := \left\{ \sum_i a_i x^i \in A[[x]] \mid a_i \pi^{ij} \rightarrow 0 \text{ for } i \rightarrow \infty \right\},$$

the algebra of power series with coefficients in A converging on a closed disk of radius $|\pi^{-j}|$. Set $A\langle \Delta^n \rangle_{\pi^j} := A\langle x_0, \dots, x_n \rangle_{\pi^j} / (x_0 + \dots + x_n - 1)$. For varying n these assemble into a simplicial ring $A\langle \Delta^\bullet \rangle_{\pi^j}$. For each j , the classifying space of the simplicial group $\text{GL}(A\langle \Delta^\bullet \rangle_{\pi^j})$ is the infinite loop space of a connective spectrum which we denote by $KV^{(j)}(A)$.

Remark. The homotopy groups of the spectrum $KV^{(0)}(A)$ are the K -groups of the Banach algebra A defined by Karoubi-Villamayor [3].

For varying j we get a pro-spectrum $KV^{\text{an}}(A) := \{KV^{(j)}(A)\}_{j \in \mathbb{N}}$ called *analytic KV-theory*. In order to globalize to all rigid spaces, we need a descent result for affinoid coverings. It turns out that one needs a non-connective variant of analytic KV -theory to achieve this. We construct this using an analog of the classical Bass construction, replacing (Laurent) polynomial rings by rings of convergent power series $A\langle x \rangle_1$ and $A\langle x, x^{-1} \rangle_1$. The resulting pro-spectrum $KH^{\text{an}}(A)$ is called *analytic KH-theory*.

3. RESULTS

In this section we assume that the characteristic of k is 0.

Theorem 1. *Let A_0 be an algebra of topologically finite type over \mathcal{O}_K and assume that $A := A_0 \otimes_{\mathcal{O}_K} K$ is regular. Then there is a homotopy cartesian square*

$$\begin{array}{ccc} K^B(A_0) & \longrightarrow & K^{B,\text{cont}}(A_0) \\ \downarrow & & \downarrow \\ K^B(A) & \longrightarrow & KH^{\text{an}}(A) \end{array}$$

in $\text{Pro}(\text{Sp}^+)$.

Here K^B is the non-connective Bass K -theory introduced by Thomason-Trobaugh [5] and $K^{B,\text{cont}}$ is defined similarly as before. $\text{Pro}(\text{Sp}^+)$ is the category of pro-objects of bounded above spectra. It is a localization of $\text{Pro}(\text{Sp})$.

Theorem 2. *The functor $KH^{\text{an}}: (\text{affinoid algebras}) \rightarrow \text{Pro}(\text{Sp}^+)$ satisfies descent for affinoid coverings.*

The second theorem allows us to extend analytic KH -theory to all rigid spaces. Theorem 1 shows that the resulting theory fits in the desired homotopy cartesian square.

The proofs of these results use pro-cdh descent [4], Weibel's K -dimension conjecture, and resolution of singularities. This is where we need the characteristic 0 assumption.

REFERENCES

- [1] S. Bloch, H. Esnault, M. Kerz, *p-adic deformation of algebraic cycle classes*, Invent. Math. 195 (2014), no. 3, 673–722.
- [2] S. Bloch, H. Esnault, M. Kerz, *Deformation of algebraic cycle classes in characteristic zero*, Algebr. Geom. 1 (2014), no. 3, 290–310.
- [3] M. Karoubi and O. Villamayor, *K-théorie algébrique et K-théorie topologique. I*, Math. Scand. 28 (1971), 265–307.
- [4] M. Morrow, *Pro cdh-descent for cyclic homology and K-theory*, J. Inst. Math. Jussieu 15 (2016), no. 3, 539–567
- [5] R. W. Thomason and T. Trobaugh, *Higher algebraic K-theory of schemes and of derived categories*, The Grothendieck Festschrift, Vol. III, 247–435, Progr. Math., 88, Birkhäuser Boston, Boston, MA, 1990.

Homotopy t -structure and effectivity

FRÉDÉRIC DÉGLISE

(joint work with Mikhail Bondarko)

1. INTRODUCTION

In this talk, I have presented a work in collaboration with Mikhail Bondarko in the context of relative motives.

Voevodsky's theory of motivic complexes, over a perfect field, first comes as a category of effective objects in which the Tate twist is non invertible for the tensor product. On the other hand, it is equipped with a canonical t -structure called the *homotopy t -structure*. This t -structure allows us to have a deeper understanding of motivic complexes, in particular in the case of curves. The t -structure can be extended to the non effective case, but then its properties are less strong; in particular, geometric motives are not bounded for sure.

The work done in collaboration with Bondarko aims at extending the notion of effectivity to the case of motives relative to an arbitrary base. Though candidates for this notion already exist, they do not satisfy good functorial properties. We define a notion of effective motives over an arbitrary base scheme S as a full subcategory of the category motives over S . The definition allows us to control the behaviour of these motives under the six operations constructed in the stable case along the lines of Ayoub.

Then we are able to extend the definition of Voevodsky's t -structure to these effective motives, and obtain an extension of certain good properties proved by Voevodsky over a field to the case of an arbitrary base. Besides, this new homotopy t -structure shares good similar properties to that of perverse sheaves, extending the work of Ayoub on the *perverse homotopy t -structure* (cf. [Ayo07]).

2. EFFECTIVE MOTIVES AND DIMENSION FUNCTIONS

For a base scheme S (excellent noetherian finite dimensional scheme), we let $DM(S)$ be the triangulated category of mixed motives with rational coefficients (cf. [CD12]) or, if S is an \mathbb{F} -scheme for \mathbb{F} a prime field of exponential characteristic p , with $\mathbb{Z}[1/p]$ -coefficients (cf. [CD15]). We will use a special motive associated with a separated S -scheme of finite type type $f : X \rightarrow S$, called the *Borel-Moore motive* of X/S :

$$M^{BM}(X/S) := f_!(\mathbb{1}_X),$$

where $\mathbb{1}_X$ is the constant motive over X .

Our main tool to define effective motives is that of a *dimension function*¹ $\delta : S \rightarrow \mathbb{Z}$ over the scheme S . Once such a dimension function is chosen, the definition goes on as follows.

¹see [ILO14, chap. 2], or [BD15, 1.1] for recall;

Definition 1. (cf. [BD15, Def. 2.2.1]) We define the category of δ -effective motives over S , denoted by $DM^{\delta\text{-eff}}(S)$, as the localizing triangulated subcategory of $DM(S)$ generated by motives of the form $M(X/S)(n)$ for any separated S -scheme X of finite type and any integer $n \geq \delta(X)$.²

Example 2. When $S = \text{Spec}(k)$ is the spectrum of a perfect field, and $\delta = 0$ is the obvious dimension function, we get that $DM^{\delta\text{-eff}}(k)$ is equivalent to Voevodsky's category of (unbounded) motivic complexes (cf. [BD15, Ex. 2.3.12]).

The category of δ -effective motives satisfies good functorial properties:

- (1) Given a closed immersion i with complementary open immersion j , it is stable under the functors i^* , $i_* = i_!$, $j_!$, j^* . In particular, the 2-functor $DM^{\delta\text{-eff}}$ satisfies the so-called *localization property* (cf. [BD15, Cor. 2.2.10]).
- (2) More generally, it is stable under the functor $f_!$ (resp. $f^*(d)$) where f is a separated morphism of finite type (resp. any morphism of finite type whose dimension of fibers is bounded by d) (cf. [BD15, Cor. 2.2.6]).

3. THE δ -HOMOTOPY t -STRUCTURE IN THE EFFECTIVE CASE

The triangulated category $DM(S)$, as well as $DM^{\delta\text{-eff}}(S)$ defined above, is compactly generated. It is well known that one can define t -structure on these type of triangulated categories \mathcal{T} by choosing homologically positive objects³, as a full sub-category of \mathcal{T} which is stable under direct sums, positive shift and extensions (see [BD15, §1.1] for recall). After Keller and Vossieck, we will call such a sub-category an *aisle* of \mathcal{T} .

Definition 3. Let S be a scheme equipped with a dimension function δ .

One defines the δ -homotopy t -structure on $DM^{\delta\text{-eff}}(S)$ as the unique one such that the (homologically) positive objects are the smallest aisle containing motives of the form: $M^{BM}(X/S)(\delta(X) + n)[2\delta(X) + n]$ for any separated S -scheme X and any integer $n \geq 0$.

Example 4. In the condition of the example in the previous section, the δ -homotopy t -structure on $DM^{\delta\text{-eff}}(k)$ coincides with Voevodsky's homotopy t -structure.

The δ -homotopy t -structure has good functorial properties; we refer the reader to the introduction of [BD15] for details. We are able to prove these properties as we can control the homology with respect to this t -structure by the following tool.

Definition 5. Let M be a δ -effective motive over S and $i \in \mathbb{Z}$ an integer. Given a couple (x, n) where $x \in S(E)$ is a point with values in a field E (finitely generated

²The notation $\delta(X)$, for any pro-étale S -scheme stands for the maximum of the integers $\text{tr.deg.}(\kappa(x)/\kappa(x))$ where x runs over the points of X and s is the projection of X to S .

³Our conventions for t -structure will be homological;

over S) and $n \leq 0$ an integer, we define the δ -homology at (x, n) with coefficients in M as follows:

$$\widehat{H}_i(M)(x, n) := \varinjlim_X [\mathrm{Hom}(M^{BM}(X/S)(\delta(X) - n)[2\delta(X) - n + i], M)]$$

where $X = \mathrm{Spec}(A)$ runs over S -models of x .⁴ We look at $\widehat{H}_i(M)$ as a functor on the discrete category whose objects are the couples (x, n) as above.

Our main theorem is the following one.

Theorem 1. ([BD15, Th. 3.3.1]) *Under the notations of the previous definition, the following conditions on a δ -effective motive M over S are equivalent:*

- (i) M is positive (resp. negative) for the δ -homotopy t -structure;
- (ii) for any integer $i \leq 0$ (resp. $i \geq 0$), $\widehat{H}_i^\delta(M) = 0$.

In other words, \widehat{H}_i^δ behaves like the homology with respect to the δ -homotopy t -structure. During the talk, we have drawn several consequences of this theorem.

- (1) *Functorial properties.*— Let $f : T \rightarrow S$ is smooth of finite type (resp. proper and with dimension of fibers bounded by d) then $f^! : DM^{\delta\text{-eff}}(T) \rightarrow DM^{\delta\text{-eff}}(S)$ is t -exact (resp. $f_* : DM^{\delta\text{-eff}}(S) \rightarrow DM^{\delta\text{-eff}}(T)$ has homological amplitude $[0, d]$) for the δ -homotopy t -structure.
- (2) *Boundedness.*— If S is regular and $\delta = -\mathrm{codim}_S$, the constant motive $\mathbb{1}_S$ in $DM^{\delta\text{-eff}}(S)$ belongs to the heart of the δ -homotopy t -structure.⁵
- (3) *Link with perversity.*— A δ -effective motive M over S is homologically positive (resp. negative) for the δ -homotopy t -structure if and only if the following condition holds:

$$\forall x \in S, H_p^{\delta_x}(i_x^! M) = 0 \text{ when } p \leq \delta(x) \text{ (resp. } p \geq \delta(x)).$$

Here δ_x is the obvious dimension function induced by δ on the spectrum of the residue field $\kappa(x)$ of x in S , and we have put $i_x^! = j^*i^!$ for the factorization $\mathrm{Spec}(\kappa(x)) \xrightarrow{j} \overline{\{x\}} \xrightarrow{i} S$.

It is striking that the last property is the exact analog of the characterization of the perverse t -structure as defined by Beilinson, Bernstein and Deligne.

REFERENCES

- [Ayo07] J. Ayoub, *Les six opérations de Grothendieck et le formalisme des cycles évanescents dans le monde motivique (I, II)*, Astérisque, vol. 314, 315, Soc. Math. France, 2007.
- [BD15] M.V. Bondarko and F. Déglise. Dimensional homotopy t -structure. arXiv: 1512.06044, 2015.
- [CD12] D.C. Cisinski and F. Déglise, *Triangulated categories of mixed motives*, arXiv:0912.2110v3, 2012.
- [CD15] ———, *Integral mixed motives in equal characteristics*, Doc. Math. (2015), no. Extra volume: Alexander S. Merkurjev's sixtieth birthday, 145–194.

⁴*i.e.* affine S -schemes of finite type such that $A \subset E$.

⁵One will be careful that this result is false if we drop the assumption that S is regular or if we consider the analog of the δ -homotopy t -structure on the stable category $DM(S)$.

[ILO14] L. Illusie, Y. Laszlo, and F. Orgogozo (eds.), *Travaux de Gabber sur l'uniformisation locale et la cohomologie étale des schémas quasi-excellents*, Astérisque, vol. 361-362, S.M.F., 2014.

Homology of GL_3 of elliptic curves

MATTHIAS WENDT

In the talk, I reported some progress in the ongoing project of constructing the elliptic dilogarithm complex using group homology. Here, *elliptic dilogarithm complex* refers to a complex computing the weight-two parts of the K-theory of an elliptic curve. Such a complex has been described by Goncharov and Levin in their work on Zagier's conjecture for elliptic curves [1]. The overall goal of the project is to identify this (essentially two-term) complex as a differential in a spectral sequence computing group homology of $\mathrm{GL}_4(k[E])$, similar to the group homology approach to the motivic weight-two complex over fields due to Suslin and Dupont–Sah. Hopefully, different realizations of the elliptic dilogarithm complex and the resulting alternative presentations for the first and second K-groups of elliptic curves can help to improve our understanding of K-theory of elliptic curves.

For the formulation of the results of the group homology computation, let k be an arbitrary field, let \overline{E}/k be an elliptic curve with fixed k -rational point P , and set $E = \overline{E} \setminus \{P\}$. The computations so far describe the homology of the group $\mathrm{GL}_3(k[E])$, using its action on the Bruhat–Tits building \mathfrak{B} associated to GL_3 and the field $k(E)$ equipped with the valuation v_P associated to the k -rational point P . The key result is the following computation of the action of $\mathrm{GL}_3(k[E])$ on the building \mathfrak{B} .

Theorem 1. (1) *The quotient $\mathrm{GL}_3(k[E]) \backslash \mathfrak{B}$ has the homotopy type $\Sigma \mathcal{F}l_3(k)$, the suspension of the flag complex for the vector space k^3 . This is essentially the union of the links of the two stable rank 3 bundles on \overline{E} appearing in the building.*

(2) *The subcomplex of cells with non-trivial stabilizer deformation-retracts equivariantly to a graph of groups Γ_E . The underlying graph of Γ_E is given by the following diagram of moduli spaces of vector bundles on \overline{E} :*

$$\mathcal{M}_{2,1}(\overline{E}) \leftarrow \mathcal{M}_{2,0}(\overline{E}) \rightarrow \mathcal{M}_{3,0}(\overline{E}).$$

Here $\mathcal{M}_{r,d}(\overline{E})$ denotes the moduli space of vector bundles of rank r and degree d on \overline{E} . However, the (semi-)stable bundles \mathcal{V} in $\mathcal{M}_{2,i}$ actually correspond to the rank 3 bundles $\mathcal{V} \oplus \det^{-1} \mathcal{V}$ on \overline{E} . This becomes a graph of groups by mapping vector bundles to their automorphism groups.

The basic proof steps are the following: first, the identification of orbits of vertices with suitable vector bundles on the curve \overline{E} is classical; this identification moreover identifies the stabilizers of vertices as automorphism groups of the corresponding bundles. Using Atiyah's classification of vector bundles on elliptic curves, explicit computations of the action of automorphism groups on the links and explicit identification of the corresponding elementary transformations

of vector bundles, one obtains a complete description of the quotient. The simplified description in the theorem then arises from an explicit deformation retraction which moves unstable bundles towards semistable bundles.

From the isotropy spectral sequence for the action of $\mathrm{GL}_3(k[E])$ on \mathfrak{B} together with classical stabilization results in group homology, we get an exact sequence, where the morphism in brackets is the group homology version of the elliptic dilogarithm complex:

$$H_2(\Gamma_E) \rightarrow K_2(E) \rightarrow \left[\mathrm{St}_3(k)/\mathcal{R} \xrightarrow{\partial} H_1(\Gamma_E) \right] \rightarrow K_1(E) \rightarrow 0.$$

Here $H_1(\Gamma_E)$ is the first homology group of the graph of groups (which is defined via the cone of the inclusion map from edge groups into vertex groups). The explicit identification of the graph Γ_E allows to identify $H_1(\Gamma_E)$ with the subspace of $\mathrm{Jac}_{\overline{E}}(\overline{k}) \otimes \overline{k}^\times$ generated by $Q \otimes u$ where Q is a point of degree ≤ 3 and $u \in k(Q)^\times$. This presentation is a lot smaller (in terms of number of generators) than the usual presentation of $K_1(\overline{E})$ as Somekawa K-group.

The Steinberg representation $\mathrm{St}_3(k)$ (which arises from the homology of the flag complex $\mathcal{F}l_3(k)$) gives rise to a geometric interpretation of K_2 -classes as triangles in \mathbb{P}^2 . The relations \mathcal{R} are not explicitly described for now; for this, computations of the action of $\mathrm{GL}_4(k[E])$ on the associated building are necessary. Part of the differential ∂ can be seen in this geometric description as mapping a side L of the triangle to the vector bundle associated to the degree 3 divisor obtained by intersecting the line L with the curve $E \subset \mathbb{P}^2$.

In future work, it remains to explicitly compute ∂ and \mathcal{R} . Naturally, the expectation would be that there is a morphism of complexes from the elliptic dilogarithm complex described by Goncharov and Levin [1] to the complex above arising from group homology. This morphism of complexes should induce the restriction map $K_i(\overline{E}) \rightarrow K_i(E)$ on homology. Note, however, that the above complex does in fact provide integral computations, where Goncharov and Levin compute rational K-theory.

Finally, as an interesting aside, the computation of the quotient $\mathrm{GL}_3(k[E]) \backslash \mathfrak{B}$ actually allows to completely determine the structure of $H^\bullet(\mathrm{GL}_3(k[E]); \mathbb{F}_\ell)$ (as module over the ring of Chern classes $\mathbb{F}_\ell[c_1, c_2, c_3]$) when \overline{E} is an elliptic curve defined over the finite field $k = \mathbb{F}_q$. When $\ell \mid q - 1$, this provides new counterexamples to a function field analogue of Quillen's conjecture on cohomology of arithmetic groups; the reason behind the counterexamples is essentially that the non-contractibility of both the quotient $\mathrm{GL}_3(k[E]) \backslash \mathfrak{B}$ as well as the graph of vector bundles with non-trivial automorphisms implies the existence of many cohomology classes which are torsion for the Chern class ring $\mathbb{F}_\ell[c_1, c_2, c_3]$.

A preliminary version of Theorem 1 (without proofs) and further discussion may be found in a note on the arXiv [2].

REFERENCES

- [1] A. Goncharov and A. Levin. *Zagier's conjecture for $L(E, 2)$* . Invent. Math. 132 (1998), 393–432.
 [2] M. Wendt, *Homology of GL_3 over function rings of elliptic curves*. arXiv:1501.02613v1.

Motives, nearby cycles and Milnor fibers

FLORIAN IVORRA

(joint work with Joseph Ayoub and Julien Sebag)

In this talk, we give a survey of the main results of [4, 8] relating tubes in non-archimedean geometry, nearby motivic sheaves and motivic nearby cycles in the sense of Denef-Loeser.

Let k be a field of characteristic zero, $R = k[[t]]$ be the ring of formal power series and $K = k((t))$ be its fraction field. In this exposition, we restrict ourselves to the stable homotopy category of schemes $\mathbf{SH}(k)$ introduced by Morel and Voevodsky and its rigid analytic version $\mathbf{RigSH}(K)$ introduced by Ayoub.

1. RECOLLECTIONS ON RIGID MOTIVES

Let X be a smooth rigid variety over K . We denote by $M_{\text{rig}}(X)$ the homological motive associated with X and by

$$M_{\text{rig}}^{\vee}(X) := \underline{\mathbf{Hom}}(M_{\text{rig}}(X), \mathbb{1}_{\text{Spm}(K)})$$

its dual. Recall that, given a separated K -scheme of finite type X , there is an associated rigid analytic K -variety X^{an} . The functor $X \mapsto X^{\text{an}}$ extends into a triangulated functor

$$\mathbf{Rig}^* : \mathbf{SH}(K) \rightarrow \mathbf{RigSH}(K)$$

such that $\mathbf{Rig}^*(M(X)) = M_{\text{rig}}(X^{\text{an}})$ where $M(X)$ is the usual (algebraic) motive of the K -scheme X .

Let $\mathbf{QUSH}(k)$ be the category of quasi-unipotent (algebraic) motives, i.e., the full triangulated subcategory of $\mathbf{SH}(\mathbf{G}_{m,k})$ closed under infinite direct sums and generated by the objects of the form $\text{Sus}_T^p(Q_r^{gm}(X, g) \otimes \mathbb{1})$ where X is a smooth k -scheme, $g \in \mathcal{O}(X)^\times$, $r \in \mathbf{N}^\times$ and $Q_r^{gm}(X, g)$ is the smooth $\mathbf{G}_{m,k}$ -scheme

$$Q_r^{gm}(X, g) := \text{Spec}(\mathcal{O}_X[T, T^{-1}, V]/(V^r - gT)) \rightarrow \text{Spec}(k[T, T^{-1}]) = \mathbf{G}_{m,k}.$$

In [3], Ayoub shows that the composition of the three functors

$$\mathbf{QUSH}(k) \hookrightarrow \mathbf{SH}(\mathbf{G}_{m,k}) \xrightarrow{t^*} \mathbf{SH}(K) \xrightarrow{\mathbf{Rig}^*} \mathbf{RigSH}(K)$$

is an equivalence of categories. We fix a quasi-inverse to the above equivalence

$$\mathfrak{R} : \mathbf{RigSH}(K) \xrightarrow{\sim} \mathbf{QUSH}(k)$$

and are interested in the composition

$$1^* \circ \mathfrak{R} : \mathbf{RigSH}(K) \rightarrow \mathbf{SH}(k)$$

where $1 : \text{Spec}(k) \rightarrow \mathbf{G}_{m,k}$ is the unit section.

2. TUBES AND NEARBY MOTIVIC SHEAVES

Let $f: X \rightarrow \text{Spec}(R)$ be a separated finite type R -scheme and denote by X_η and X_σ the generic and special fibers of X .

By t -adic completion, one obtains a separated formal R -scheme topologically of finite type $\mathcal{X} \rightarrow \text{Spf}(R)$ and we denote by \mathcal{X}_η the generic fiber of \mathcal{X} . The rigid analytic variety \mathcal{X}_η is an open analytic subvariety of the analytification X_η^{an} of the algebraic generic fiber X_η . With a locally closed subset $Z \subset X_\sigma$, we may associate its tube $]Z[$ which is an open rigid analytic subvariety of \mathcal{X}_η .

Assume that the rigid analytic variety \mathcal{X}_η is smooth over K . One of the main result of [4] is the following theorem that relates rigid motives of tubes with the nearby motivic sheaf $\Psi_f(\mathbf{1}_{X_\eta})$ introduced by Ayoub in [2].

Theorem 1 (see [4]). *Denote by $z : Z \hookrightarrow X_\sigma$ the inclusion. Then, there is a canonical isomorphism*

$$1^* \circ \mathfrak{R}(M_{\text{rig}}^\vee(]Z[)) \simeq (f_\sigma)_* z_* z^* \Psi_f(\mathbf{1}_{X_\eta})$$

in the category of motives $\mathbf{SH}(k)$.

Taking $Z = X_\sigma$, one gets that the cohomological motive $M_{\text{rig}}^\vee(\mathcal{X}_\eta)$ is related to the nearby motivic sheaf by a canonical isomorphism

$$1^* \circ \mathfrak{R}(M_{\text{rig}}^\vee(\mathcal{X}_\eta)) \simeq (f_\sigma)_* \Psi_f(\mathbf{1}_{X_\eta})$$

in $\mathbf{SH}(k)$.

Our main theorem is a motivic analog of a theorem of Berkovich that we recall now. Let \overline{K} be the completion of an algebraic closure of the valued field K and let \overline{k} be its residue field. Set $\overline{Z} = Z \times_k \overline{k}$ and $]\overline{Z}[=]Z[\widehat{\times}_K \overline{K}$. In [5, 6], Berkovich constructed a canonical isomorphism of étale cohomology groups

$$\mathbb{H}_{\text{ét}}^i(]\overline{Z}[, \mathbf{Q}_\ell) \simeq \mathbb{H}_{\text{ét}}^i(\overline{Z}, R\Psi_f(\mathbf{Q}_{\ell, X_\eta})|_{\overline{Z}}).$$

It is worth noting that Berkovich’s theorem holds over general non-archimedean fields whereas, to state Theorem 1, we need to assume equal characteristic zero. Indeed, this is required in [3] to ensure the existence of the equivalence \mathfrak{R} .

3. RELATION WITH THE WORK OF DENEFF AND LOESER

Let X be a semi-stable R -scheme. Denote by $(D_i)_{i \in I}$ be the irreducible components of its special fiber and set

$$D_J := \bigcap_{j \in J} D_j \quad D_J^\circ := D_J \setminus \bigcup_{i \notin J} D_i$$

for a non-empty subset $J \subseteq I$.

Theorem 2 (see [4, 8]). *Let X be a semi-stable R -scheme. For $\emptyset \neq J \subset I$, let $\rho_J : \widetilde{D}_J^\circ \rightarrow D_J^\circ$ be the étale finite cover defined as in [8, §3.1.3]. Then, one has the formula*

$$[\Psi_f(\mathbf{1}_{X_\eta})] = \sum_{\emptyset \neq J \subset I} (-1)^{|J|-1} \left[M_{X_\sigma, c}(\widetilde{D}_J^\circ \times_k \mathbf{G}_{m, k}^{|J|-1}) \right]$$

in $K_0(\mathbf{SH}_{\text{ct}}(X_\sigma))$.

Note that $\mathbf{SH}_{\text{ct}}(X_\sigma)$ is the full triangulated subcategory of $\mathbf{SH}(X_\sigma)$ formed by the constructible motives. In [7], Denef and Loeser have introduced a notion *motivic nearby cycles* which is an element ψ_f in the Grothendieck ring of the special fiber \mathcal{M}_{X_σ} and is defined in terms of the arc scheme as the limit of the motivic zeta function associated with f .

Corollary 1 (see [4, 8]). *Let X be a finite type R -scheme with smooth generic fiber and denote by $f : X \rightarrow \text{Spec}(R)$ its structural morphism. We have the equality*

$$[\Psi_f(\mathbf{1}_{X_\eta})] = \chi_{X_\sigma, c}(\psi_f)$$

in $K_0(\mathbf{SH}_{\text{ct}}(X_\sigma))$.

Here

$$\chi_{X_\sigma, c} : \mathcal{M}_{X_\sigma} \rightarrow K_0(\mathbf{SH}_{\text{ct}}(X_\sigma))$$

is the motivic Euler characteristic over the special fiber X_σ . A particular case of Theorem 1 gives an isomorphism of motives

$$1^* \circ \mathfrak{R}(\mathbf{M}_{\text{rig}}^\vee(\mathcal{F}_x)) \simeq x^* \Psi_f(\mathbf{1}_{X_\eta})$$

where \mathcal{F}_x is the analytic Milnor fiber at x introduced by Nicaise-Sebag in [9]. From this and Corollary 1, we deduce the following formula

$$[1^* \circ \mathfrak{R}(\mathbf{M}_{\text{rig}}^\vee(\mathcal{F}_x))] = \chi_{k, c}(\psi_{f, x})$$

where $\psi_{f, x} := x^* \psi_f$ in \mathcal{M}_k and $\chi_{k, c} : \mathcal{M}_k \rightarrow K_0(\mathbf{SH}_{\text{ct}}(k))$ is the motivic Euler characteristic over k . This formula expresses the fact that the motivic Milnor fiber $\psi_{f, x}$ of Denef-Loeser, at least as a class in the Grothendieck ring of constructible motives, is determined by the rigid motive of the analytic Milnor fiber.

REFERENCES

- [1] J. Ayoub, *Les six opérations de Grothendieck et le formalisme des cycles évanescents dans le monde motivique. I*, Astérisque **314** (2007).
- [2] J. Ayoub, *Les six opérations de Grothendieck et le formalisme des cycles évanescents dans le monde motivique. II*, Astérisque **315** (2007).
- [3] J. Ayoub, *Motifs des variétés analytiques rigides*, Mém. Soc. Math. Fr. (N.S.) **140–141** (2015).
- [4] J. Ayoub, F. Ivorra, J. Sebag, *Motives of rigid analytic tubes and nearby motivic sheaves*, Submitted for publication, (2014).
- [5] V. Berkovich, *Vanishing cycles for formal schemes*, Invent. Math. **115** (1994), no 3, 539–571.
- [6] V. Berkovich, *Vanishing cycles for formal schemes. II*, Invent. Math. **125** (1996), no 2, 367–390.
- [7] J. Denef, F. Loeser, *Motivic Igusa zeta functions*, J. Algebraic Geom. **7** (1998), 505–537.
- [8] F. Ivorra, J. Sebag, *Nearby motives and motivic nearby cycles*, Selecta Math. (N.S.) **19** (2013), no 4, 879–902.
- [9] J. Nicaise, J. Sebag, *Motivic Serre invariants, ramification, and the analytic Milnor fiber*, Invent. Math. **168** (2007), no 1, 133–173.

A motivic version of Segal's theorem

IVAN PANIN

(joint work with G.Garkusha)

Using the machinery of framed sheaves developed by Voevodsky [Voe2] we state and prove a motivic Segal's theorem. It states particularly that for any infinite perfect field k and any integer $n > 0$ a natural morphism

$$\underline{Hom}(\Delta^\bullet \times -, \Omega_{\mathbb{P}^1}^\infty \Sigma_T^\infty(T^n)) \rightarrow \Omega_{\mathbb{P}^1, mot}^\infty \Sigma_T^\infty(T^n)$$

of motivic spaces is a stalk-wise weak equivalence for the Nisnevich topology. Here $T^n = \mathbb{A}^n / (\mathbb{A}^n - \{0\})$ is the motivic $(2n, n)$ -sphere, $\Omega_{\mathbb{P}^1}$ is the naive \mathbb{P}^1 -loop-functor and $\Omega_{\mathbb{P}^1, mot}$ is the motivic \mathbb{P}^1 -loop-functor. To prove this result a triangulated category of framed motives $SH_{S^1}^{fr}(k)$ is introduced and studied.

We also construct a compactly generated triangulated category of framed bispectra $SH^{fr}(k)$. Using an extension of the above theorem it is proved that $SH^{fr}(k)$ reconstructs the Voevodsky category $SH(k)$ in the case of an infinite and perfect field.

As a topological application, it is proved that in the complex number case for any integer $n > 0$ there is an equality in the ordinary homotopy category of simplicial sets

$$Fr(\Delta_{\mathbb{C}}^\bullet, T^n) = \Omega_{S^1}^\infty \Sigma_{S^1}^\infty(S^{2n}),$$

where S^{2n} is the ordinary $2n$ -sphere.

REFERENCES

- [AGP] *Ananyevskiy, A., Garkusha, G., Panin, I.* Cancellation theorem for framed motives of algebraic varieties, preprint arXiv:1601.06642.
- [BF] *Bousfield, A.K., E. M. Friedlander, E.M.* Homotopy theory of Γ -spaces, spectra, and bisimplicial sets, In Geometric applications of homotopy theory II, Lecture Notes in Math. 658, Springer Verlag, 1978, pp. 80-130.
- [GP1] *Garkusha, G., Panin, I.* Framed motives of algebraic varieties (after V. Voevodsky), preprint arXiv:1409.4372.
- [GP2] *Garkusha, G., Panin, I.* Homotopy invariant presheaves with framed transfers, preprint, 2015, arXiv:1504.00884.
- [GP3] *Garkusha, G., Panin, I.* Linear framed motives of algebraic varieties, preprint, 2016 (in preparation).
- [GNP] *Garkusha, G., Neshitov, A., Panin, I.* Framed motives of relative motivic spheres, arXiv:1604.02732v2
- [MV] *Morel, F., Voevodsky, V.* F. Morel, V. Voevodsky, A^1 -homotopy theory of schemes, Publ. Math. IHES 90 (1999), 45-143.
- [S] *Segal, G.* Categories and cohomology theories, Topology 13 (1974), 293-312.
- [Voe1] *Voevodsky, V.* Triangulated category of motives over a field, in Cycles, Transfers and Motivic Homology Theories (V. Voevodsky, A. Suslin and E. Friedlander, eds.), Annals of Math. Studies, Princeton University Press, 2000.
- [Voe2] *Voevodsky, V.* Notes on framed correspondences, unpublished, 2001-2003.
- [Voe3] *Voevodsky, V.* A^1 -Homotopy theory, Doc. Math., Extra Vol. ICM 1998(1), 417-442.

Euler class groups and the homology of elementary groups

MARCO SCHLICHTING

A ring R has many units if for every integer $n \geq 1$, there are central elements $a_1, \dots, a_n \in R$ such that for all $\emptyset \neq I \subset \{1, \dots, n\}$, the partial sum $a_I = \sum_{i \in I} a_i$ is a unit in R . For instance, any infinite field, any commutative local ring with infinite residue field, any (possibly non-commutative) algebra over a ring with many units has many units. For a ring R , denote by $E_n(R)$ the subgroup of $GL_n(R)$ generated by the elementary matrices. For a commutative ring R and integer $n \geq 1$, we let $SL_n(R)$ denote the subgroup of $GL_n(R)$ consisting of matrices of determinant one (for the definition of SL_0 ; see below). For a discrete group G , we denote by $H_n(G)$ its integral homology groups, that is, $\text{Tor}_n^{\mathbb{Z}G}(\mathbb{Z}, \mathbb{Z})$. Finally, denote by $sr(R) \in \mathbb{N}$ the stable range of R . For instance, $sr(R) = 1$ for any commutative local ring, $sr(R) = 1$ for any (possibly non-commutative) artinian ring, and $sr(R) \leq 1 + \dim R$ when R is commutative noetherian.

Theorem 1 ([Sch]). *Let R be ring with many units. Then*

- (1) $H_i(E_n R, E_{n-1} R) = 0, \quad i \leq n - sr(R)$
- (2) *If R is commutative, then*

$$H_i(SL_n R, SL_{n-1} R) = 0, \quad i \leq n - sr(R).$$

When R is a field of characteristic zero, Theorem 1 was previously proved by Hutchinson and Tao [HT10]. The theorem implies a conjecture of Bass [Bas73].

Theorem 2 ([Sch]). *Let R be a ring with many units. Then*

$$\pi_i(BGL_n^+ R, BGL_{n-1}^+) = 0 \quad i \leq n - sr(R).$$

Bass actually conjectured Theorem 2 for commutative noetherian rings without the hypothesis of "many units". However, in this general case, $R = \mathbb{Z}$ already provides a counter example.

Let A be a commutative ring, let $\mathbb{Z}[A^*]$ be the group ring of the group of units A^* in A with standard \mathbb{Z} -basis $\langle a \rangle, a \in A^*$, multiplication $\langle a \rangle \cdot \langle b \rangle = \langle ab \rangle$, and $\langle 1 \rangle = 1$. Let $I_{A^*} = \text{Ker}(\mathbb{Z}[A^*] \rightarrow \mathbb{Z} : \langle a \rangle \mapsto 1)$ be the augmentation ideal, and let $[a] = \langle a \rangle - 1 \in I_{A^*}$.

Definition 3 ([Sch]). *Let A be a commutative ring (with infinite residue fields). We define the graded $\mathbb{Z}[A^*]$ -algebra*

$$\widehat{K}_*^{MW}(A) = \text{Tens}_{\mathbb{Z}[A^*]} I_{A^*} / \text{Steinberg}$$

as the quotient of the tensor algebra of I_{A^} over the group ring $\mathbb{Z}[A^*]$ modulo the ideal generated by the Steinberg relations $[a] \otimes [1 - a]$ for $a, 1 - a \in A^*$.*

For instance,

$$\widehat{K}_0^{MW}(A) = \mathbb{Z}[A^*], \quad \widehat{K}_1^{MW}(A) = I_{A^*}, \quad \widehat{K}_2^{MW}(A) = I_{A^*} \otimes_{A^*} I_{A^*} / \text{Steinberg}.$$

We prove in [Sch] that the group $\widehat{K}_n^{MW}(A)$ is isomorphic to the Morel-Hopkins Milnor-Witt K -group $K_n^{MW}(A)$ when $n \geq 2$ and A is a field or local ring with

residue field $\neq \mathbb{F}_2, \mathbb{F}_3$. We have the following strengthening of Theorem 1. Here, SL_0A is defined to be the discrete set of units A^* in A , so as to make the formula $H_i(SL_nA) = \text{Tor}_i^{\mathbb{Z}[GL_nA]}(\mathbb{Z}[A^*], \mathbb{Z})$ hold for all i, n .

Theorem 4 ([Sch]). *Let A be a commutative local ring with infinite residue field. Then there are isomorphisms of A^* -modules for all $n \geq 0$*

$$H_i(SL_nA, SL_{n-1}A; \mathbb{Z}) \cong \begin{cases} 0 & i < n \\ \widehat{K}_n^{MW}(A) & i = n. \end{cases}$$

This theorem is the SL_n -analogue of a theorem of Nesterenko-Suslin [NS89]. A version of the theorem was proved for fields of characteristic zero by Hutchinson-Tao [HT10].

Let R be a noetherian ring with infinite residue fields, and let P be an oriented projective R -module of rank n . We define the ‘‘Euler class’’ $e(P)$ of P as a certain Zariski cohomology class

$$e(P) \in H_{Zar}^n(R, \mathcal{K}_n^{MW})$$

where \mathcal{K}_n^{MW} denotes the Zariski sheaf associated with the presheaf $A \mapsto \widehat{K}_n^{MW}(A)$. Using the previous theorem, we can show the following.

Theorem 5 ([Sch]). *Let R be a commutative noetherian ring of dimension $n \geq 2$. Assume that all residue fields of R are infinite. Let P be an oriented rank n projective R -module. Then*

$$P \cong Q \oplus R \Leftrightarrow e(P) = 0 \in H_{Zar}^n(R, \mathcal{K}_n^{MW}).$$

Using \mathbb{A}^1 -homotopy theory, a version of this theorem was proved by Morel [Mor12] for smooth algebras over infinite perfect fields.

REFERENCES

- [Bas73] Hyman Bass. Some problems in ‘‘classical’’ algebraic K -theory. In *Algebraic K-theory, II: ‘‘Classical’’ algebraic K-theory and connections with arithmetic (Proc. Conf., Battelle Memorial Inst., Seattle, Wash., 1972)*, pages 3–73. Lecture Notes in Math., Vol. 342. Springer, Berlin, 1973.
- [HT10] Kevin Hutchinson and Liqun Tao. Homology stability for the special linear group of a field and Milnor-Witt K -theory. *Doc. Math.*, (Extra volume: Andrei A. Suslin sixtieth birthday):267–315, 2010 .
- [Mor12] Fabien Morel. \mathbb{A}^1 -algebraic topology over a field, volume 2052 of *Lecture Notes in Mathematics*. Springer, Heidelberg, 2012.
- [NS89] Yu. P. Nesterenko and A. A. Suslin. Homology of the general linear group over a local ring, and Milnor’s K -theory. *Izv. Akad. Nauk SSSR Ser. Mat.*, 53(1):121–146, 1989.
- [Sch] Marco Schlichting. Euler class groups, and the homology of elementary and special linear groups. arXiv:1502.05424v3.

Participants

Prof. Dr. Alexey Ananyevskiy

Chebyshev Laboratory
St. Petersburg State University
14th Line 29B
Vasilyevsky Island
199 178 St. Petersburg
RUSSIAN FEDERATION

Dr. Giuseppe Ancona

Institut für Mathematik
Universität Zürich
Winterthurerstrasse 190
8057 Zürich
SWITZERLAND

Dr. Aravind Asok

Department of Mathematics
University of Southern California
3620 South Vermont Ave., KAP 108
Los Angeles, CA 90089-2532
UNITED STATES

Dr. Christian Ausoni

Département de Mathématiques
Institut Galilée
Université Paris XIII
99 av. J.B. Clément
93430 Villetaneuse Cedex
FRANCE

Dr. Federico Binda

Fachbereich Mathematik
Universität Duisburg-Essen
45117 Essen
GERMANY

Dr. Oliver Bräunling

Mathematisches Institut
Universität Freiburg
Eckerstrasse 1
79104 Freiburg i. Br.
GERMANY

Prof. Dr. Ulrich Bunke

Fakultät für Mathematik
Universität Regensburg
93040 Regensburg
GERMANY

Prof. Dr. Guillermo Cortinas

Dep. de Matemática/IMAS
Facultad de Ciencias
Exactas y Naturales
Universidad de Buenos Aires
Ciudad Universitaria Pab. 1
C 1428 EGA Buenos Aires
ARGENTINA

Dr. Frederic Deglise

Mathématiques
École Normale Supérieure de Lyon
46, Allée d'Italie
69364 Lyon Cedex 07
FRANCE

Dr. Veronika Ertl

Fakultät für Mathematik
Universität Regensburg
Universitätsstrasse 31
93053 Regensburg
GERMANY

Prof. Dr. Dr. h.c.mult. Hélène Esnault

FB Mathematik und Informatik
Freie Universität Berlin
Arnimallee 3
14195 Berlin
GERMANY

Prof. Dr. Jean Fasel

Institut Fourier
UMR 5582, CNRS/UJF
Université de Grenoble I
100 rue de Maths
P.O. Box 74
38402 Saint-Martin-d'Hères
FRANCE

Prof. Dr. Thomas Geisser

Department of Mathematics
Rikkyo University
Nishi-Ikebukuro
Tokyo 171
JAPAN

Prof. Dr. Christian Haesemeyer

Department of Mathematics
University of California, Los Angeles
Box 951555
Los Angeles CA 90095-1555
UNITED STATES

Daniel Harrer

Mathematisches Institut
Universität Freiburg
Eckerstrasse 1
79104 Freiburg i. Br.
GERMANY

Prof. Dr. Lars Hesselholt

Graduate School of Mathematics
Nagoya University
Chikusa-ku, Furo-cho
Nagoya 464-8602
JAPAN

Prof. Dr. Jens Hornbostel

FB C: Mathematik u.
Naturwissenschaften
Bergische Universität Wuppertal
Gaußstrasse 20
42119 Wuppertal
GERMANY

Dr. Marc Hoyois

Department of Mathematics
Massachusetts Institute of Technology
77 Massachusetts Avenue
Cambridge, MA 02139-4307
UNITED STATES

Dr. Florian Ivorra

U. F. R. Mathématiques
I. R. M. A. R.
Université de Rennes I
Campus de Beaulieu
35042 Rennes Cedex
FRANCE

Prof. Dr. Uwe Jannsen

Fakultät für Mathematik
Universität Regensburg
93040 Regensburg
GERMANY

Prof. Dr. Bruno Kahn

Institut de Mathématiques
Université Paris VI
Case 247
4, Place Jussieu
75252 Paris Cedex 05
FRANCE

Dr. Wataru Kai

Graduate School of Mathematical
Sciences
University of Tokyo
3-8-1 Komaba, Meguro-ku
Tokyo 153-8914
JAPAN

Dr. Shane Kelly

Mathematisches Institut
Albert-Ludwigs-Universität Freiburg
Eckerstraße 1
79104 Freiburg i. Br.
GERMANY

Prof. Dr. Moritz Kerz

Fakultät für Mathematik
Universität Regensburg
Universitätsstrasse 31
93053 Regensburg
GERMANY

Dr. Tohru Kohrita

Graduate School of Mathematics
Nagoya University
Chikusa-ku, Furo-cho
Nagoya 464-8602
JAPAN

Prof. Dr. Marc Levine

Fakultät für Mathematik
Universität Duisburg-Essen
Universitätsstrasse 2
45117 Essen
GERMANY

Prof. Dr. Stephen Lichtenbaum

Department of Mathematics
Brown University
Box 1917
Providence, RI 02912
UNITED STATES

Prof. Dr. Alexander S. Merkurjev

Department of Mathematics
University of California at
Los Angeles
405 Hilgard Avenue
Los Angeles, CA 90095-1555
UNITED STATES

Dr. Baptiste Morin

Institut de Mathématiques de Toulouse
Université Paul Sabatier
31062 Toulouse Cedex 9
FRANCE

Dr. Matthew Morrow

Hausdorff Center for Mathematics
Endenicher Allee 60
53115 Bonn
GERMANY

Prof. Dr. Ivan A. Panin

Laboratory of Algebra at St. Petersburg
Department of V.A. Steklov Institute of
Mathematics (P.O.M.I.)
Russian Academy of Sciences
27, Fontanka
191 023 St. Petersburg
RUSSIAN FEDERATION

Dr. Simon Pepin-Lehalleur

Institut für Mathematik
Freie Universität Berlin
Arnimallee 6
14195 Berlin
GERMANY

Prof. Dr. Gereon Quick

Institutt for Matematiske fag
NTNU Trondheim
7491 Trondheim
NORWAY

Prof. Dr. Oliver Röndigs

Fachbereich Mathematik/Informatik
Universität Osnabrück
49069 Osnabrück
GERMANY

Prof. Dr. Andreas Rosenschon

Mathematisches Institut
Ludwig-Maximilians-Universität
München
Theresienstrasse 39
80333 München
GERMANY

Dr. Kay Rülling

FB Mathematik und Informatik
Freie Universität Berlin
Arnimallee 3
14195 Berlin
GERMANY

Dr. Kanetomo Sato

Department of Mathematics
Chuo University
1-13-27, Kasuga, Bunkyo-ku
Tokyo 112-8552
JAPAN

Dr. Marco Schlichting

Mathematics Institute
University of Warwick
Zeeman Building
Coventry CV4 7AL
UNITED KINGDOM

Prof. Dr. Alexander Schmidt

Mathematisches Institut
Universität Heidelberg
Im Neuenheimer Feld 205
69120 Heidelberg
GERMANY

Jun. Prof. Dr. Markus Spitzweck

Fachbereich Mathematik/Informatik
Universität Osnabrück
49069 Osnabrück
GERMANY

Prof. Dr. Vasudevan Srinivas

School of Mathematics
Tata Institute of Fundamental Research
Homi Bhabha Road
Mumbai 400 005
INDIA

Dr. Rin Sugiyama

Department of Mathematics
Tokyo Denki University
Senjuasahicho 5, Adachi-ku
Tokyo 120-8551
JAPAN

Dr. Goncalo Tabuada

Department of Mathematics
Massachusetts Institute of Technology
Cambridge, MA 02139-4307
UNITED STATES

Dr. Georg Tamme

Fakultät für Mathematik
Universität Regensburg
93040 Regensburg
GERMANY

Prof. Dr. Charles Vial

Department of Pure Mathematics
and Mathematical Statistics
University of Cambridge
Wilberforce Road
Cambridge CB3 0WB
UNITED KINGDOM

Konrad Völkel

Mathematisches Institut
Universität Freiburg
Eckerstrasse 1
79104 Freiburg i. Br.
GERMANY

Prof. Dr. Mark E. Walker

Department of Mathematics
University of Nebraska, Lincoln
Lincoln NE 68588
UNITED STATES

Prof. Dr. Charles A. Weibel

Department of Mathematics
Rutgers University
Busch Campus, Hill Center
New Brunswick, NJ 08854-8019
UNITED STATES

Dr. Matthias Wendt

Fakultät für Mathematik und Physik
Leibniz-Universität Hannover
Welfengarten 1
30167 Hannover
GERMANY

Dr. Yigeng Zhao

Fakultät für Mathematik
Universität Regensburg
Universitätsstrasse 31
93053 Regensburg
GERMANY

Dr. Takao Yamazaki

Mathematical Institute
Faculty of Science
Tohoku University
6-3 Aoba
Sendai 980-8578
JAPAN

Dr. Changlong Zhong

Department of Mathematics
State University of New York at Albany
1400 Washington Avenue
Albany, NY 12222
UNITED STATES

Dr. Inna Zakharevich

Department of Mathematics
Cornell University
320 Tower Road
Ithaca, NY 14850
UNITED STATES

